

Skull Sexual Monomorphism in the Gwembe Dwarf Goat Breed from Zambia

Parés-Casanova PM¹, Kataba A², Mwaanga ES² and Simukoko H²

¹Department of Animal Production, University of Lleida, Av. Alcalde Rovira Roure, Lleida, Catalunya, Spain

²Department of Biomedical Sciences, University of Zambia, School of Veterinary Medicine, Lusaka, Zambia

*Corresponding author: Parés-Casanova PM, Department of Animal Production, University of Lleida, Av. Alcalde Rovira Roure 191, E-25198 Lleida, Catalunya, Spain, E-mail: peremiquelp@prodan.udl.cat

Citation: Parés-Casanova PM, Kataba A, Mwaanga ES, Simukoko H (2014) Skull Sexual Monorphism in the Gwembe Dwarf Goat Breed from Zambia. J Vet Sci Anim Husb 1(5): 503

Abstract

This study involved some morphometric parameters of the skull of thirty adult Gwembe Dwarf Goats (15 males and 15 females) without any apparent skeletal disorders. Lower jaws were not included in this study. A total of 43 linear measurements were analyzed. The analysis reflected no differences between sexes, thereby indicating lack of general conformation differences between males and females. However, comparison of the parameters that exhibited most variance revealed that horn core basal circumference and length of the horn core on the front margin were strongly influenced by sex. Therefore, only the horn conformation could be used as discriminator sexual variable. This result couples with the fact that males usually have stronger and larger horns than females but skull size seeming no to be very different between sexes. The obtained results would indicate a marked sexual monomorphism in this breed.

Keywords: Capra hircus; Craniometry; Sexual dimorphism; Skull biometry

Introduction

Local goat has great genetic resource potential that can be utilized as a source of superior breeding formulation adaptable to local conditions. [1] Reported that breeds of local livestock are important and should be protected because of their ability to survive under low-quality feed, unfavourable local climatic conditions, and their high resistance to local diseases and parasites. These abilities are critical in cross-breeding ventures aimed at sustaining genetic pools that have evolved through centuries within the given locality.

Phenotypic and morphological characteristics are still commonly used by researchers and practitioners in characterization of animals' breeds [2]. The Gwembe Dwarf Goat or locally known as "Mpongo" is a small breed with average weight of 35 kg for both males and females found in the Gwembe valley in Southern Province of Zambia. Well-adapted to hot and dry climatic conditions with low rainfall patterns, this breed is of multi colour coat variations ranging from completely black, brown, black and white, grey to white and brown- with horns of medium size and usually curved backwards (<http://dad.fao.org/>). Other goat breeds of Zambia include the Plateau Goat located on the plateau regions of Southern Province, the Sinazongwe district of the Southern and North part of the Gwembe Valley, and the imported Boer and Saanen (<http://dad.fao.org/>). The morphologic and morphometric studies of the head region do not only reflect contributions of genetic and environmental components to individual development, but describe genetic and ecophenotypic variation [3]. Until now, there have been numerous comparative morphological studies of the skull anatomy in many of the mammalian species. In particular, in small ruminants the morphological structures and geometrical measurements of the skull bones have been examined to detect the distinguishing features of these species [4] and breeds [3,5,6]. To the best knowledge of the authors, there is no published data on the morphometric parameters of the head region applied anatomy in the Zambian native goats. An exception of this is the thesis by [7], which does not contain detailed information on head conformation other than the head length and width for male and female goats of unknown breed. Moreover, complete comparative studies between sexes are not available.

So, the morphometric values of the skull of the Gwembe Dwarf Goat will not only be of ethnological importance but they will undoubtedly aid to better the description of sexual differences. Furthermore, the obtained results will provide important baseline data for further comparative studies on the skull of the Zambian native goat breeds.

Materials and Methods

Thirty dry skulls of adult Gwembe Dwarf Goat specimens aged 18 months and older were collected from five different local farms and studied. The age was determined on the basis of the growth of the third molar tooth [8]. All specimens used for this study showed no cranial deformities or bony injuries. The total series included all specimens collected.

The method described by [9] was used for assessing cranial (neuro- and viscerocranium) linear measurements, and it is commonly used in other osteometric studies [10]. A total of 43 linear measurements were analyzed (Table 1). Lower jaws were not included in this study. Basic descriptive statistics were calculated for all samples and no size adjustment was applied. The measurements were performed using 0.1 mm callipers and an ovinometer. All measurements were taken by the same person (Kataba).

Parameter	Definition
1	Total length of the skull: the distance between akrokranion-prosthion
2	Condylbasal length: caudal border of occipital condyles-prosthion
3	Total length of the cranial base: basion-prosthion
4	Short skull length: basion-premolare
5	Premolare-prosthion length
7	Upper length of the viscerocranium: nasion-prosthion
8	Median frontal length: akrokranion-nasion
9	Akrokranion-bregma length
10	Frontal length: bregma-nasion
11	Upper neurocranium length: akrokranion-supraorbitale
12	Facial length: supraorbitale-prosthion
13	Akrokranion-infraorbitale length
14	Greatest length of the lacrimal bone: most lateral point of the lacrimal bone –most rostral point of the lacrimomaxillary suture
15	Greatest length of the nasal bone: nasion-rhinion
16	Entorbitale-prosthion length
17	Distance between the caudal border of one occipital condyle and the infraorbitale of the same side
18	Dental length: postdentale-prosthion
19	Oral palatal length: palatinoorale-prosthion
20	Lateral length of the premaxilla: nasointermaxillare-prosthion
21	Length of the maxillary tooth row
22	Length of the upper molar row
23	Length of the upper premolar row
24	Greatest inner width of the orbit: ectorbitale-entorbitale
25	Greatest inner height of the orbit
26	Greatest mastoid breadth: otion-otion
27	Greatest breadth of the occipital condyles
28	Greatest breadth at the bases of the paracondylar processes
29	Greatest breadth of the foramen magnum
30	Height of the foramen magnum: basion-opisthion
31	Least breadth of parietal
32	Greatest breadth between the lateral borders of horn core base
33	Greatest neurocranium breadth: euryon-euryon
34	Greatest frontal breadth: ectorbitale-ectorbitale
35	Least breadth between the orbits: entorbitale-entorbitale
36	Facial breadth: between facial tuberosities
37	Greatest breadth across the nasal bones
38	Greatest breadth across the premaxilla
39	Greatest palatal breadth
40	Horncore basal circumference
41	Greatest diameter of horn core base
42	Least diameter of horn core base
43	Length of the horn core on the front margin
44	Distance between horn tips

Table 1: Studied parameters (nomenclature according to Von den Driesch, 1976). A total of 43 linear measurements were analyzed. Von den Driesch's numeration has been served, except for var. 44 which is ours.

Main simple descriptive statistics (range, mean, standard deviation and coefficient of variation) were obtained, as well the to assess normality of distribution of data in both sexes. The non-parametric one-way NPMANOVA test (Non-Parametric MANOVA, also known as PERMANOVA) was applied, using the Mahalanobis distance [11]. The significance was computed by permutation of group membership, with 9,999 replicates. Given the large number of parameters, a selection of the most relevant parameters was ultimately made using a multivariate analysis of principal components (PCA). The principal components (PC) were extracted from the co-variance matrix. The eigenanalysis was carried out on the group means ("between-group", i.e. the items analysed are the groups, not the specimens). The loadings within the individual eigenvectors for each PC were used to form contrasts in order to describe the skulls numerically. A one-way MANOVA (Multivariate ANalysis Of VAriance) tested whether male and female samples had the same mean for the most discriminant parameters. The two-sample Hotelling's T^2 test was avoided as some of these parameter were non parametric. In order to identify and measure the associations among the two sets of parameters, a Canonical Correspondence Analysis was done, according to [12] and the two-tailed non-parametric U Mann-Whitney was used to test whether the medians of samples (males and females) were different.

The data recorded were analyzed statistically in customary fashion using PAST® (v. 2.17c) software. Level confidence was established at 5%.

Results

In Table 2 there appear the main descriptive statistics for linear measurements. The NPMANOVA reflected no differences between sexes ($F=1.104$, $p=0.317$), so it can be supposed that there appeared no general conformation differences between males and females.

Var.	♀							♂						
	Min	Max	X	StD	CV	W	P	Min	Max	X	StD	CV	W	P
1	13.0	16.7	15.1	1.319	8.7	0.908	0.125	13.4	16.7	15.0	0.858	5.7	0.944	0.438
2	12.6	16.4	14.3	1.389	9.7	0.897	0.085	12.5	15.8	14.7	0.746	5.1	0.801	0.004
3	11.8	14.5	13.1	0.786	6.0	0.969	0.839	11.6	14.1	12.6	0.757	6.0	0.953	0.573
4	7.7	10.3	9.1	0.789	8.7	0.963	0.742	7.5	10.3	8.2	0.685	8.3	0.783	0.002
5	3.5	4.6	4.0	0.348	8.6	0.956	0.622	3.5	4.3	4.0	0.254	6.4	0.951	0.541
7	7.8	9.8	8.6	0.669	7.8	0.921	0.198	7.5	9.4	8.5	0.534	6.3	0.974	0.917
8	7.7	10.0	9.1	0.680	7.5	0.946	0.464	7.8	9.8	8.6	0.536	6.2	0.939	0.365
9	4.6	5.3	5.0	0.197	3.9	0.951	0.533	4.3	5.2	4.9	0.246	5.0	0.924	0.218
10	5.3	6.4	5.9	0.340	5.8	0.967	0.815	5.3	6.6	6.0	0.369	6.1	0.947	0.484
11	8.2	9.3	8.9	0.342	3.8	0.888	0.063	8.0	10.0	9.1	0.711	7.8	0.928	0.251
12	9.0	10.6	9.8	0.403	4.1	0.981	0.976	8.4	10.3	9.5	0.563	5.9	0.93	0.269
13	9.9	12.0	10.6	0.615	5.8	0.923	0.213	10.0	11.5	10.7	0.443	4.1	0.931	0.285
14	1.9	3.3	2.6	0.369	14.2	0.991	1.000	2.0	2.9	2.6	0.218	8.5	0.948	0.491
15	3.9	5.6	4.8	0.556	11.7	0.943	0.422	4.0	5.5	4.6	0.411	9.0	0.889	0.065
16	7.5	9.9	8.8	0.739	8.4	0.966	0.791	7.7	9.7	8.9	0.578	6.5	0.959	0.671
17	10.1	11.9	11.0	0.658	6.0	0.928	0.251	9.9	11.8	11.1	0.525	4.7	0.94	0.383
18	6.9	8.7	7.9	0.588	7.4	0.923	0.212	6.6	8.5	7.6	0.467	6.1	0.981	0.976
19	5.7	6.8	6.2	0.371	5.9	0.854	0.020	5.1	6.5	6.0	0.377	6.3	0.947	0.484
20	5.1	6.3	5.6	0.439	7.8	0.849	0.017	4.7	6.0	5.5	0.401	7.3	0.936	0.333
21	4.3	5.6	5.0	0.449	9.0	0.918	0.180	4.1	5.3	4.6	0.388	8.5	0.887	0.060
22	2.5	3.1	2.8	0.149	5.2	0.926	0.237	2.6	2.9	2.8	0.097	3.5	0.903	0.107
23	1.5	2.7	2.2	0.443	20.5	0.873	0.038	1.4	2.5	1.9	0.403	20.7	0.887	0.060
24	2.8	3.3	3.1	0.171	5.5	0.952	0.556	2.8	3.4	3.1	0.188	6.0	0.934	0.314
25	2.9	3.2	3.1	0.093	3.0	0.944	0.429	2.8	3.4	3.2	0.167	5.3	0.981	0.973
26	5.4	6.2	5.9	0.223	3.8	0.884	0.055	5.2	6.2	5.8	0.308	5.3	0.924	0.218
27	3.8	4.4	4.1	0.168	4.1	0.942	0.402	4.0	4.3	4.1	0.087	2.1	0.92	0.192
28	4.8	5.9	5.3	0.251	4.7	0.954	0.589	5.0	5.7	5.3	0.177	3.3	0.95	0.522
29	1.6	2.0	1.9	0.106	5.7	0.973	0.897	1.5	2.0	1.8	0.136	7.5	0.913	0.151
30	1.5	1.8	1.6	0.093	5.7	0.93	0.269	1.4	1.7	1.6	0.105	6.6	0.936	0.331
31	2.2	3.9	2.9	0.430	14.7	0.952	0.556	2.7	3.7	3.1	0.318	10.1	0.946	0.461

Var.	♀							♂						
	Min	Max	X	StD	CV	W	P	Min	Max	X	StD	CV	W	P
32	5.5	7.2	6.4	0.416	6.5	0.981	0.976	6.2	7.1	6.8	0.276	4.1	0.928	0.256
33	5.6	6.6	5.9	0.264	4.5	0.795	0.003	5.4	6.0	5.8	0.191	3.3	0.95	0.524
34	7.3	9.1	8.2	0.485	5.9	0.936	0.331	7.6	8.5	8.1	0.283	3.5	0.942	0.413
35	5.4	7.3	6.4	0.479	7.5	0.975	0.922	5.7	6.9	6.3	0.272	4.3	0.949	0.511
36	5.0	6.1	5.6	0.333	5.9	0.938	0.357	5.1	6.0	5.6	0.244	4.4	0.982	0.979
37	1.6	2.5	2.1	0.237	11.1	0.983	0.984	1.7	2.4	2.1	0.232	11.2	0.944	0.433
38	1.7	2.3	2.1	0.189	9.0	0.905	0.112	1.7	2.3	2.1	0.151	7.1	0.961	0.716
39	5.0	5.9	5.5	0.275	5.0	0.963	0.747	5.0	5.9	5.4	0.246	4.5	0.979	0.959
40	0.0	8.0	6.4	1.954	30.5	0.672	0.000	6.8	9.0	8.2	0.615	7.5	0.94	0.382
41	1.8	2.9	2.4	0.328	13.8	0.962	0.732	2.2	3.2	2.8	0.260	9.3	0.982	0.982
42	1.5	2.1	1.9	0.208	11.2	0.897	0.085	1.9	2.5	2.1	0.169	7.9	0.905	0.114
43	0.0	6.8	4.9	1.611	33.1	0.817	0.006	4.5	8.4	6.0	1.089	18.2	0.928	0.251
44	4.5	7.9	5.9	0.939	15.9	0.968	0.830	6.0	8.9	7.0	0.748	10.7	0.881	0.048

X: average
 StD: Standard Deviation
 CV: Coefficient of Variation (%)
 W: Shapiro-Wilk

Table 2: Main descriptive statistics for linear measurements (expressed in mm) for females (♀, n=15) and males (♂, n=15). Von den Driesch's numeration has been served, except for var. 44 which is ours. Not normally distributed data appear in bold.

For the PCA, first 6 PCs presented a eigenvalue higher of Jolliffe cut-off's one (0.363). PC1 explained a 43% of variance (Table 3). The most important parameters that entered into the PC1 were 40 (horncore basal circumference) and 43 (length of the horn core on the front margin), both related to horns, and into the PC2 was 40. For PC3 was the parameter 4 (short skull length), which presented the highest value (Table 4). For these parameters, NPMANOVA did reflect differences between sexes (F=4.752, p=0.006), so sexual dimorphism can be suspected not for a general skull form but only for horns. Horn parameters that had strong influence on the sex were precisely the length of the horn core on the front margin (1.239, Can1) and the horncore basal circumference (-0.935, Can1) (Figure 1), the latter one (but not the former) (U=65.5, p=0.053) being statistically different between sexes (U=20.5, p<0.001) (Figure 2).




Figure 1: Canonical Analysis for horn parameters (95% confidence ellipses), showing the position of individuals according to their horn conformation (squares for males, n=15, crosses for females, n=15). Number on dots show each analyzed parameter (42: least diameter of horn core base; 43: length of the horn core on the front margin; 44: distance between horn tips).




Figure 2: Histogram of horncores basal circumferences for males (right, n=15) and females (left, n=15). This parameter that shows statistically differences between sexes (Mann-Whitney's $U=20.5$, $p<0.001$).

Pc	Eigenvalue	% variance	% total explained
1	6.486	43.042	43.042
2	3.422	22.705	65.747
3	1.369	9.085	74.832
4	0.925	6.141	80.973
5	0.691	4.586	85.559
6	0.390	2.589	88.148

Table 3: Eigenvalues and % variances for first 6 Principal Components (PCs). Jolliffe cut-off=0.363.

	PC1	PC2	PC3
1	0.263	0.359	-0.330
2	0.337	0.282	-0.325
3	0.110	0.295	0.335
4	0.012	0.281	0.498
5	0.048	0.108	-0.080
7	0.106	0.143	0.161
8	-0.051	0.133	0.331
9	0.018	0.033	0.034
10	0.064	0.034	-0.092
11	0.042	-0.056	0.301
12	0.061	0.111	0.171
13	0.145	0.091	-0.097
14	0.064	0.090	0.007
15	0.093	0.127	0.005
16	0.144	0.153	-0.189
17	0.181	0.116	0.031
18	0.124	0.214	0.027

	PC1	PC2	PC3
19	0.069	0.164	0.013
20	0.094	0.121	0.038
21	0.062	0.142	0.091
22	-0.001	0.027	-0.002
23	0.062	0.112	0.004
24	0.033	0.040	-0.065
25	0.015	-0.004	0.069
26	0.023	0.061	0.027
27	0.013	-0.002	-0.010
28	0.035	0.051	-0.001
29	-0.016	0.002	-0.002
30	-0.017	0.002	-0.011
31	-0.043	-0.098	-0.131
32	0.090	-0.050	0.032
33	0.023	0.038	-0.031
34	0.045	0.130	0.011
35	0.082	0.125	0.049
36	0.057	0.094	-0.004
37	0.013	0.039	0.037
38	0.025	0.025	0.003
39	0.055	0.083	0.053
40	0.541	-0.439	0.091
41	0.072	-0.113	0.132
42	0.043	-0.047	-0.038
43	0.506	-0.252	0.092
44	0.264	-0.161	0.159

Table 4: Component loading for each parameter for first 3 Principal Component (PC), which explained the 75% of the total observed variance. Highest absolute values for each component appear in bold.

Discussion

Dimorphism means two forms “sexual dimorphism” means that the two sexes of a species differ in external appearance or other features. Males and females may differ in size, color, shape, the development of appendages (such as horns, teeth, feathers, or fins), and also in scent or sound production. Species in which male and females are identical in appearance or other features are said to be “monomorphic”. Our results showed that there are no general skull conformation differences between males and females in the Dwarf Zambian Goat, although some particular parameters have an strong influence on the breed sex, such as the length of the horn core on the front margin, the distance between horn tips, and the short skull length. This is in line with the fact that males usually have stronger and larger horns than females [8]. Short skull length (parameter 4) has low influence, so size seems not to be very different between sexes. According to [7] 2009, males and females would differ by their head width, not by their head length, but, as earlier stated, his research is not clearly focused on Dwarf goat.

The global variation between sexes may be interpreted in relation to extensive management styles of the animals that are moreover under a low anthropogenic influence so they tend to reinforce the natural sexual size dimorphism of the species. Thus, horn conformation could be used as discriminator sexual variable and this fact can be attributed to that sexual dimorphism in this breed is not related to differences in the diet of the two sexes but probably to participation in fighting between males, or to a mere artificial selection of males towards bigger horns by farmers.

References

1. Food and Agriculture Organization (2007) Global Plan of Action for Animal Genetic resources and Interlaken Declaration on Animal Genetic Resources, Rome, Italy.
2. Batubara A, Noor RR, Farajallah A, Tiesnamurti B, Doloksaribu M (2011) Morphometric and phylogenic analysis of six population Indonesian local goats. J Ani Sci Tech 34: 165-74.
3. Monfared AL, Naji H, Sheibani MT (2013) Applied Anatomy of the Head Region of the Iranian Native Goats (*Capra Hircus*). Global Vet 10: 60-4.

4. Onuk B, Kabak M, Atalar K (2013) Anatomic and craniometric factors in differentiating roe deer (*Capreolus capreolus*) from sheep (*Ovis aries*) and goat (*Capra hircus*) skulls. *Arch Biol Sci* 65: 133-41.
5. Sarma K (2006) Morphological and Craniometrical Studies on the Skull of Kagani Goat (*Capra hircus*) of Jammu Region. *Int J Morphol* 24: 449-55.
6. Shawulu JC, Kwari HD, Olopade JO (2011) Morphology of the bones of the skull in the Sahel ecotypes of goats (*Capra hircus*) in Nigeria. *J Vet Anat* 4: 1-13.
7. Ahmadu B (2001) Management Practices, Morphometric Body Measurements and Diseases of Local Zambian Goats. Ph. Thesis. The University of Zambia, Lusaka, Zambia.
8. Dyce KM, Sack WO, Wensing CJG (2002) *Textbook of Veterinary Anatomy* (3rd edn) Saunders Publishers, Philadelphia, USA.
9. Driesch A Von den (1976) *A Guide to the Measurement of Animal Bones from Archaeological sites*. Peabody Museum Press, Massachusetts, USA.
10. Onar V, Belli O, Owen PR (2005) Morphometric Examination of Red Fox (*Vulpes vulpes*) from the Van-Yoncatepe Necropolis in Eastern Anatolia. *Int J Morphol* 23: 253-60.
11. Anderson MJ (2001) A new method for non-parametric multivariate analysis of variance. *Austral Ecology* 26: 32-46.
12. Legendre P, Legendre L (1998) *Numerical Ecology* (2nd edn) Elsevier Publications, Oxford, UK.