Universitat de Lleida

Document downloaded from:

<http://hdl.handle.net/10459.1/68009>

The final publication is available at:

<https://doi.org/10.1007/s12550-019-00362-y>

Copyright

(c) Society for Mycotoxin (Research Gesellschaft für Mykotoxinforschung e.V.)
and Springer-Verlag GmbH Germany, part of Springer Nature, 2019

Mycotoxin Research

Influence of the ingredients used in the formulation of dairy cattle feedstuffs on aflatoxin contamination of feed and milk

--Manuscript Draft--

Manuscript Number:	MYRE-D-19-00004
Full Title:	Influence of the ingredients used in the formulation of dairy cattle feedstuffs on aflatoxin contamination of feed and milk
Article Type:	Original Article
Abstract:	<p>Aflatoxins (AFs) are mycotoxins produced by <i>Aspergillus</i> species that may contaminate a wide variety of feedstuffs. Aflatoxin M1 (AFM1) is a hydroxylated metabolite of aflatoxin B1 (AFB1) and can be excreted in milk of cattle after consuming aflatoxin-contaminated feed. The aim of this research was to assess the level of AFs in dairy cattle feedstuffs (193 samples) and the level of AFM1 in raw milk samples (375) estimating the AFB1/AFM1 carry-over from feed to milk. Moreover, the correlation between the raw materials used as ingredients of the total mixed ration (TMR) and the presence of AFs was studied. 34.7% of the feed samples analysed presented AFs in a range of 0.05-6.45 µg/kg, being 12.4% positive for AFB1. 18.9% of the milk samples contained AFM1 with concentrations ranging from 0.009 to 1.36 µg/kg. None of the feed samples exceeded the EU legal limit of AFB1, while three raw milk samples presented levels of contamination higher than the maximum level established by the EU. A carry-over rate from 0.6 to 6% was estimated. The statistical analysis showed that maize silage, bagasse, soya bean husk, corn, alfalfa hay, cotton seed and compound feed were normally presented as ingredients of AFB1-positive feed samples, thus special attention should be paid in controlling these raw materials when used in TMR preparation. Few studies have focussed before in the impact of TMR components in the AFs contamination of the final feedstuff.</p>

Influence of the ingredients used in the formulation of dairy cattle feedstuffs on aflatoxin contamination of feed and milk

M. Rodríguez-Blanco, A.J. Ramos, M. Prim, V. Sanchis and S. Marín

Applied mycology unit, Food technology department, University of Lleida, UTPV-XaRTA, Agrotecnio Centre, Av. Rovira Roure 191, 25198 Lleida, Spain

*Corresponding author

Sonia Marín Sillué

Tel.: +34 973702542

E-mail: smarin@tecal.udl.cat

ORCID

M. Rodríguez-Blanco 0000-0003-1667-8222

A.J. Ramos 0000-0002-2830-8299

V. Sanchis 0000-0001-9889-8098

S. Marín 0000-0002-0714-6155

Abstract

Aflatoxins (AFs) are mycotoxins produced by *Aspergillus* species that may contaminate a wide variety of feedstuffs. Aflatoxin M₁ (AFM₁) is a hydroxylated metabolite of aflatoxin B₁ (AFB₁) and can be excreted in milk of cattle after consuming aflatoxin-contaminated feed. The aim of this research was to assess the level of AFs in dairy cattle feedstuffs (193 samples) and the level of AFM₁ in raw milk samples (375) estimating the AFB₁/AFM₁ carry-over from feed to milk. Moreover, the correlation between the raw materials used as ingredients of the total mixed ration (TMR) and the presence of AFs was studied. 34.7% of the feed samples analysed presented AFs in a range of 0.05-6.45 µg/kg, being 12.4% positive for AFB₁. 18.9% of the milk samples contained AFM₁ with concentrations ranging from 0.009 to 1.36 µg/kg. None of the feed samples exceeded the EU legal limit of AFB₁, while three raw milk samples presented levels of contamination higher than the maximum level established by the EU. A carry-over rate from 0.6 to 6% was estimated. The statistical analysis showed that maize silage, bagasse, soya bean husk, corn, alfalfa hay, cotton seed and compound feed were normally presented as ingredients of AFB₁-positive feed samples, thus special attention should be paid in controlling these raw materials when used in TMR preparation. Few studies have focussed before in the impact of TMR components in the AFs contamination of the final feedstuff.

Keywords

aflatoxin B₁, aflatoxin M₁, carry over, total mixed ration, raw milk

Abbreviations

AFs	Aflatoxins
AFB ₁	Aflatoxin B ₁
AFB ₂	Aflatoxin B ₂
AFG ₁	Aflatoxin G ₁
AFG ₂	Aflatoxin G ₂
AFM ₁	Aflatoxin M ₁
UHPLC-FLD	Ultra-High Performance Liquid Chromatography with Fluorescence Detector
TMR	Total Mixed Ration

Introduction

Mycotoxins are secondary metabolites produced mainly by fungi belonging to the genera *Fusarium*, *Aspergillus*, *Penicillium* and *Alternaria*. In animals, the ingestion of these fungal compounds has been associated with performance problems, diseases or even death, while, in humans, the chronic intake of low levels of mycotoxins is related to different kinds of pathologies, being cancer the most important (Fink-Gremmels 1999). Fungal growth and mycotoxins production can take place in several crops and, as a result, they can reach the food and feed chain being consumed directly by humans or thorough ingestion of food of animal origin, fed with contaminated feed.

The most common mycotoxins that may occur in feedstuffs are aflatoxins (AFs), fumonisins (FBs), zearalenone (ZEN), ochratoxin A and trichothecenes (Driehuis et al. 2008). AFs are mycotoxins produced by *Aspergillus* species, mainly by *A. flavus* and *A. parasiticus*. As a consequence of fungal contamination before or after harvest, these toxins may be found in products such as nuts, dried fruits, cereals, spices, seeds, cocoa or coffee (Juan et al. 2007; Garrido et al. 2012). Up until now, more than 20 different types of AFs have been described; however, the most important in foods are aflatoxin B₁ (AFB₁), aflatoxin B₂ (AFB₂), aflatoxin G₁ (AFG₁), aflatoxin G₂ (AFG₂) and aflatoxin M₁ (AFM₁).

AFs are highly toxic, carcinogenic, mutagenic and teratogenic compounds (Bakirdere et al. 2012). AFB₁ is the most toxic AF and it is considered as the most powerful natural hepatocarcinogenic agent in mammals. The International Agency for Research on Cancer (IARC) has classified this toxin as a Group 1 human carcinogen (IARC 2012).

AFM₁ is a hydroxylated metabolite of AFB₁ and can be excreted in milk of cattle that have been fed with aflatoxin-contaminated feed (Fallah et al. 2009; Iqbal et al. 2015). AFM₁ is not as toxic as AFB₁ and has been classified by IARC as Group 2B as possibly carcinogenic to humans (IARC 1993). AFM₁ can be detected in milk of animals that have consumed feedstuff with AFB₁ within 24-48 h and it almost disappears after 72 h (Martins and Martins 2000; Battacone et al. 2003). The transformation of AFB₁ into AFM₁ and its presence in the secreted milk depends on several factors related with feed, metabolism, weather and geographical location of dairy farms (Masoero et al. 2007; Iqbal et al. 2013). The extent of carry-over is influenced by various nutritional and physiological factors, including regimes, rate of ingestion, health of the animal, hepatic biotransformation capacity, and milk production. Consequently, the rate of absorption of AFs and the excretion of aflatoxins varies between animal, from day to day, and also from one milking to the next. It has been reported that the predicted

rate of AFB₁/AFM₁ carry-over from feed to milk is approximately 0.3-6% (Heshmati and Milani 2010). Due to its high toxicity, legal limits of AFB₁ in animal feed have been established by authorities. Directive 2002/32/EC sets an upper limit of 5.0 µg/kg in complete feedstuffs for dairy cattle with a moisture content of 12% (EC 2002). In addition, in order to reduce human exposure to AFM₁, the European Commission Regulation 1881/2006 sets a maximum permissible limit of 0.05 µg/kg for AFM₁ in raw milk, heat-treated milk and milk for the manufacture of milk-based products (EC 2006). Precisely because of the small amounts of toxin allowed, it is necessary to develop sensitive and specific analytical methods which allow the detection of very small amounts of AFs in animal feed, as well as their quantification.

Nowadays, dairy cows are usually fed using the total mixed ration (TMR) method, that consists on the supply of forages together with different kinds of concentrates or by-products (such as whole cottonseed or compound feed), grains, protein supplements, minerals and vitamins in specific quantities which are mixed thoroughly making up a balanced ration.

TMR used to feed dairy cattle is usually enriched with concentrates to achieve a high milk yield. Concentrates are low-fibre, high-energy material used as components of the ration for dairy cattle in order to raise the energy level and to increase the level of proteins, minerals (macro and micro) and fat-soluble vitamins, compensating any other deficiencies of the total ration derived from the use of the forage portion. These concentrates may be an important source of mycotoxins, as well as other materials such as cereal grains or soybean products, press cakes from oil plants (Nawaz et al. 1997; Scudamore and Livesey 1998; Placinta et al. 1999), and preserved feeding stuffs like silage, hay or straw (O'Brien et al. 2005; Mansfield and Kuldau 2007).

The aims of this work were to evaluate the occurrence of AFB₁, AFB₂, AFG₁ and AFG₂ in different dairy cattle TMR samples from farms located in four Spanish areas during 2016-2018 using an in-house validated method. Moreover, the level of AFM₁ in milk samples from cows fed with the mentioned feed was assessed in order to calculate the AFB₁/AFM₁ carry-over from feed to raw milk. The possible relation between the TMR composition and the presence of AFs in feed and, consequently, in milk, was also studied.

Materials and methods

Sampling

From February 2016 to January 2018, a total of 193 different feedstuff samples and 375 milk samples were collected from dairy farms located in different areas of Spain (Cantabria, Castilla-León, Cataluña and Galicia).

Farms with previous problems related to AFs, due to weak food safety management systems, were specifically selected. Each farm provided a bulk sample of 4-5 kg of TMR in order to have a representative fraction of the whole. The bulk sample was pooled from smaller samples collected through the day from a single batch while it was consumed by the cattle. Farmers were asked to provide a report containing the composition of each TMR sample and also the composition of the compound feed used as an additional TMR ingredient. The most common ingredients of the feed samples are shown in table 1. In addition, 37 raw materials used as TMR components including cotton seed (n=12), corn (n=6), compound feed (n=5), soya bean (n=2), okara (n=2), maize silage (n=2), dried mixture (n=1), barley silage (n=1), ray grass silage (n=1), sugar beet pulp silage (n=1), alfalfa silage (n=1), immature corn silage (n=1), sugar beet pulp (n=1) and soya bean husk (n=1), were received and they were analysed separately. Raw milk samples were collected in the dairy farms from the cows that had been fed with the sampled feed 24 hours and 48 hours before. All samples were stored at -18 °C until analysis.

Reagents and solutions

The analytical standards of AFB₁, AFB₂, AFG₁, AFG₂ and AFM₁ were supplied by Sigma (Sigma-Aldrich, Alcobendas, Spain). Acetonitrile and methanol were both HPLC grade and were obtained from Scharlab (Sentmenat, Spain). Acetic acid glacial was supplied by Fisher Scientific (Loughborough, UK). Filter paper (Whatman No. 113) was purchased from Whatman (Maidstone, UK). Immunoaffinity chromatography (IAC) columns for AFB₁, AFB₂, AFG₁ and AFG₂ (Easi-extract® Aflatoxin) and for AFM₁ (Aflaprep® M Wide) were purchased from R-Biopharm (Rhône LTD Glasgow, UK). Pure water was obtained from a Milli-Q system (Millipore, Billerica, MA, USA). Phosphate buffer saline (PBS) was prepared with potassium chloride (0.2 g) (Panreac, Castellar del Vallès, Spain), potassium dihydrogen phosphate (0.2 g) (Panreac), disodium phosphate anhydrous (1.16 g) (Panreac) and sodium chloride (8.0 g) (Panreac) in 1 L of pure water; the pH was brought to 7.4.

Preparation of stock standard solutions

Standard solutions of AFB₁, AFB₂, AFG₁ and AFG₂ were dissolved in methanol at a concentration of 250 mg/L and stored at 4 °C in a sealed vial until use. The concentrations in the stock solutions were checked by UV spectroscopy according to AOAC Official Methods of Analysis, chapter 49 (Horwitz and Latimer 2006). Working standard solutions (0.1-20 µg/L for AFB₁ and AFG₁, and 0.05-10 µg/L for AFB₂ and AFG₂) were prepared by appropriate dilution of known volumes of the stock solution with methanol:H₂O (50:50, v/v) and used to obtain calibration curves in the chromatographic system. The standard of AFM₁ was dissolved in

acetonitrile at a concentration of 1.0 mg/L and stored at 4 °C in a sealed vial until use. Working standards (0.1, 0.25, 0.5, 1, 2, 5, 10, 25, 50 µg/L) were also prepared by appropriate dilution of known volumes of the stock solution with mobile phase and used to obtain calibration curves in the chromatographic system.

Feed samples analysis

Before the analysis, the samples were dried during 24 hours at 60 °C and ground into fine powder. TMR samples were extracted and cleaned up according to the Easi-extract® Aflatoxin manual slightly modified in order to obtain higher recoveries. Briefly, five grams of feed sample were extracted with 40 mL of acetonitrile:H₂O solution (90:10, v/v) and kept in an ultrasonic bath for 10 min. The sample then was centrifuged for 10 min at 4,676 g. Three mL of supernatant were diluted with 72 mL of PBS and drained through the IAC column. The column was washed with 20 mL of PBS and AFs were eluted by applying 1 mL of methanol grade-HPLC and 1 mL of milli-Q water, consecutively. Before the injection in the chromatographic system, the final extract was filtered through a 0.22 µm PTFE disposable syringe filter (Kinesis, Cambridgeshire, UK).

The equipment used for the detection of AFs was an Agilent 1260 Infinity Quaternary LC system (Agilent Technologies, Santa Clara, California, United States) with a quaternary pump, an auto sampler, a vacuum degasser and a fluorescence detector. The chromatographic separation was carried out in a Poroshell 120 EC-C18 UHPLC column (2.7 µm particle size, 4.6 x 50 mm; Agilent Technologies) protected with a Poroshell 120 EC-C18 UHPLC Guard 3PK (2.7 µm particle size, 4.6 x 5 mm; Agilent Technologies). In order to enhance and confirm AFB₁ and AFG₁ detection, a post-column derivatization with a LCTech UVE photochemical system (LCTech GmbH, Obertaufkirchen, Germany) was performed. The fluorescence detector was set at wavelengths of 365 nm and 440 nm for excitation and emission, respectively. The mobile phase consisted of acetonitrile:methanol:water (10:20:70, v/v) and the flow rate was 1.2 mL/min. The temperature of the column was set at 40 °C, and the injection volume was 50 µL.

Milk samples analysis

Milk analysis was carried out with immunoaffinity columns, according to the ISO official method (ISO 2007). Milk samples were placed into 50-mL centrifuge tubes and were warmed to 37 °C for 30 minutes. Then, they were centrifuged for 10 min at 4,534 g. The upper cream layer was removed and the defatted supernatant was filtered with No. 113 Whatman filter papers. 40 mL of this fraction were passed through the IAC Aflaprep® M Wide column. After this, the column was washed with 20 mL of PBS and, finally, eluted twice with 1.25 mL of acetonitrile:methanol (60:40, v/v). The eluent was dried under nitrogen stream and then it was resuspended with

300 µL of mobile phase consisting of acetonitrile:methanol:0.1% acetic acid (5:15:80, v/v). Before the injection in the chromatographic system, the final extract was filtered through a 0.22 µm PTFE disposable syringe filter. The equipment used for the detection of AFM₁ was an Agilent 1260 Infinity Quaternary LC system (Agilent Technologies) with a quaternary pump, an auto sampler, a vacuum degasser and a fluorescence detector. The chromatographic separation was carried out in a Poroshell 120 EC-C18 UHPLC column (2.7 µm particle size, 4.6 x 50 mm; Agilent Technologies) protected with a Poroshell 120 EC-C18 UHPLC Guard 3PK (2.7 µm particle size, 4.6 x 5 mm; Agilent Technologies) at a temperature of 40 °C. Excitation and emission wavelengths were set at 360 nm and 450 nm, respectively. The mobile phase consisted of acetonitrile:methanol:0.1% acetic acid (5:15:80, v/v) and the flow rate was 0.8 mL/min. The injection volume was 50 µL.

Statistical analysis

The analysis of the data was performed using JMP Pro 13. Firstly, ANOVA was used so as to know if there were significant effects of year, farm and composition of TMR samples on aflatoxin levels in both feed and milk samples. A multivariate partial least square regression (PLS) was carried out in order to determine which components of the feedstuffs could explain the presence of AFs in feed. VIP (Variable Importance in Projection) scores were also calculated, which are a measurement of a variable's importance in the PLS model. They summarize the contribution of a variable to the model; higher VIP scores allow to identify the most contributory variables in class discrimination in the PLS model. Finally, correlation analysis was carried out in order to study the relationship between AFB₁ presence in feed and AFM₁ in milk.

Results

AFs occurrence in feed and milk samples

From February 2016 to January 2018, a total of 193 different TMR samples and 375 milk samples were analysed. 67 feed samples were contaminated with AFs and 71 milk samples were positive for the presence of AFM₁. A summary of the occurrence and the range of concentrations of AFs in feed and AFM₁ in milk is given in Tables 2 and 3.

In total, 24 feed samples contained AFB₁, 9 were positive for AFB₂, 47 for AFG₁ and 13 for AFG₂. Therefore, AFB₁ and AFG₁ were more frequently detected than AFB₂ and AFG₂. The concentration of AFB₁ was always lower than the EU limit of 5.0 µg/kg for complete feedstuffs for dairy cattle and only one AFG₁-positive sample had a toxin concentration that could be considered reasonably high (6.45 µg/kg). The number of AFB₁-positive

samples was higher in 2016, while in 2017 was AFG₁, the most frequent toxin found. Moreover, there seemed not to be significant differences among the four different locations under study (data not shown).

Regarding ingredients that were studied separately, AFs were not found in any of the dried mixture, maize silage, barley silage, sugar beet pulp silage, alfalfa silage, immature corn silage, sugar beet pulp, okara and soya bean husk samples. On the other hand, positive results for the presence of AFs were obtained from the analysis of corn, cotton seed, compound feed, soya bean and ray grass silage samples (Table 4).

AFM₁ was present in 18.4% of raw milk samples. 19.02% and 18.32% of the samples obtained 24 h and 48 h after feeding, respectively, presented AFM₁, therefore significant differences were not found between them.

Figure 1 shows the distribution of AFM₁ during the length of the study. The number of positive samples was clearly higher in 2016 than in 2017 and the season when more positive results were obtained was from May to July in 2016. Three raw milk samples exceeded the EU limit of 0.05 µg/kg.

Relationship between TMR composition and AFs occurrence in feed and milk samples

PLS multivariate regressions were carried out in order to assess the impact of the TMR components in AFs, AFB₁ and AFG₁ presence (Figure 2). The first model explained 49% of variability, the second 64% and the third one 37%.

AFs-positive samples usually contained maize silage, barley silage, bagasse, okara, dehydrated ray grass, straw and corn. Moreover, it has to be pointed out that the compound feed was related to the presence of group B aflatoxins in feed samples (Figure 2A). AFB₁-positive feed samples presented maize silage, bagasse, soya bean husk, corn, alfalfa hay, cotton seed and compound feed as ingredients (Figure 2B). As for AFG₁, maize silage, alfalfa silage, barley silage, okara, soya bean husk, straw, dehydrated alfalfa and cotton seed were positively related to the presence of the toxin (Figure 2C).

Due to the weight of the compound feed in the presence of AFB₁ in feed samples, a PLS analysis was carried out taking into account the components of the compound feed separately, instead of considering it as a unique element. It was not possible to get from all farmers the complete information of each TMR sample (including compound feed composition), for this reason, the analysis was performed considering only those samples whose information was available (n=33). In this subset of samples, barley silage, soya bean, corn, barley and alfalfa hay were the ingredients present in AFB₁-positive samples. They also contained corn, cotton seed and wheat as ingredients coming from the compound feed (Figure 3A). With this PLS model a 91% of the variability was explained.

As far as AFM₁ is concerned, milk positive samples were obtained from cows that had been fed with feed that usually contained ray-grass silage, wheat silage, barley silage, bagasse, dehydrated ray-grass, straw, corn, alfalfa hay and cotton seed. Also, corn, barley and cotton seed as part of the compound feed (Figure 3B). 74% of the variability was explained by the PLS model, in this case.

Carry-over from feed to milk

A total of fifteen AFM₁-positive milk samples came from cows that had been fed with AFB₁-positive feed. However, eight feed samples contained AFB₁, but milk samples obtained from cows fed with these feed turned out to be negative for the presence of AFM₁. Conversely, 27 AFM₁-positive samples came from cows fed with feed samples that were not contaminated by AFB₁. As a result, the correlation coefficient found between AFB₁ and AFM₁ was 0.4774 in milk samples obtained 24 hours after having been fed with feed samples, and 0.4529 in those samples obtained after 48 hours. Taking into account the cases in which AFs were present in both feed and milk, the daily feed intake and the milk yield by each cow, it was established that the carry-over of AFB₁ in feed into AFM₁ in milk varied from 0.6 to 6%.

Discussion

Incidence of AFs in feed

The presence of AFB₁ in animal feeds has been legislated in different countries in order to avoid the possible toxic effects related with its ingestion both in animals and in humans. The EU has set an upper limit of 5.0 µg/kg in complete feedstuffs for dairy cattle with a moisture content of 12% (EC 2002). However other countries are less restrictive and have established higher limits, such as USA, which have set an action level of AFs of 20.0 µg/kg in feeds and ingredients for dairy animals (FDA 1996).

The number of AF-contaminated samples found in this study was low, with only one sample containing a relatively high level of AFG₁ (Table 2). Our results show that 34.7% of the samples studied were contaminated by AFs in a range of 0.05-6.45 µg/kg. The most frequently found toxins were AFB₁ (12.4%) and AFG₁ (24.4%). In agreement with our research, similar results have been obtained, although it is worth noting that the farms included in our study were selected specifically due to their previous problems with AFs. Many studies found AFs in dairy feedstuffs, although only a small proportion of the contaminated samples exceeded the maximum tolerable EU limit of 5.0 µg/kg. Also in Spain, Hernández-Martínez and Navarro-Blasco, (2015) found that 90% of the dairy cow feedstuff samples under study showed detectable concentrations of AFs, without exceeding the

legal limit. In China, Han et al. (2013) obtained 53.5% AFs-positive feed samples, being 17.5% positive for AFB₁ with concentrations from 0.05 to 3.53 µg/kg. Decastelli et al. (2007) in Italy and Sassahara et al. (2005) in Brazil, detected also a low number of contaminated samples, 8.1% and 17%, respectively. In Turkey, Sahin et al. (2016) found that 26.3% of the dairy cattle feedstuffs contained AF and the concentrations ranged from 0.278 to 8.43 µg/kg, although only two samples presented levels higher than 5.0 µg/kg.

By contrast, other studies found a relatively low number of contaminated feed samples but some of them presented high values of AFs, widely exceeding the established legal limit. In Portugal, Martins et al. (2007) carried out a 10-year study finding 37.4% positive samples with a level of contamination ranging from 1-74 µg/kg. 6.2%, showed levels of AFB₁ above the EU limit (from 5.1 to 74 µg/kg). Similarly, Pleadin et al. (2015) obtained 22.2% AFB₁-positive samples and 12.3% had concentrations over the limit, showing, occasionally, very high values being the maximum found 304.6 µg/kg.

Conversely, other studies have found a higher number of samples containing AFs and in higher concentrations in dairy cattle feed. A research carried out in Turkey (Kocasari et al. 2012), observed that AFs were presented in 61.7% of samples in concentrations ranging from 3.82 to 54.34 µg/kg. In Iran, Ehsani et al. (2016) obtained 41.9% of contaminated samples with concentrations between 1.87 and 19.41 µg/kg and 42% contained AFB₁ at the level of 10-15 µg/kg. All the samples examined by Gizachew et al. (2016) in Ethiopia presented AFB₁ and 26.2% exceeded such a high concentration as 100 µg/kg. Recently, Ismail et al. (Pakistan, 2017) and Mohammed et al. (Tanzania, 2016) found 30.5% and 61.5% of the samples, respectively, with concentration of AFB₁ over the EU permissible limit.

AFM₁ in milk

Although the problem of AFM₁ in milk has been known for a long time, recently several cases of milk contaminated with this toxin in Europe have revealed the extent of the problem, both from the point of view of Public Health and from an economic angle. Thus, for example, in a sporadic outbreak that took place in the South of Spain, two million litres of raw milk, prior to its treatment and commercial sale, had to be destroyed in 2013 due to its contamination by AFM₁ above the levels legally allowed in the EU, with a cost of more than 760,000 euros (El Mundo, 2013). It cannot be ruled out that the emergence of outbreaks of contamination by AFM₁ in milk could be due, among other possible causes, to climate change, which may be causing a change in the pattern of distribution of *Aspergillus* aflatoxigenic strains in Europe and favouring their growth.

As well as with AFB₁ in animal feed, there is current regulation for the maximum allowable concentration of AFM₁ in milk in order to reduce the exposure to this toxic metabolite. The EU has set a maximum of 0.05 µg/kg while in the USA it is allowed to reach levels of AFM₁ up to 0.5 µg/kg (FDA 1996). These standards have been adopted by numerous countries (FAO 2004; EC 2006).

The present study analysed raw milk samples obtained after 24 and 48 hours, respectively, from cows that had been fed with TMR samples. 71 out of 375 milk samples (18.9%) contained AFM₁ in concentrations above the limit of detection in a range from 0.009 to 1.36 µg/kg, with only three samples exceeding the EU limit.

These results are in line with data obtained by Han et al. (2013) who observed that 22.5% of raw milk samples from China contained AFM₁ ranging from 0.005 to 0.06 µg/kg. Seven samples contained levels of toxin over the EU legal limit. Sahin et al. (2016) in Turkey, found out 21.1% of contaminated raw milk samples in a range of 0.011-0.1 µg/kg, containing three samples concentrations of AFM₁ that exceeded the EU limit. Golge (2014), also in Turkey, detected AFM₁ in 30.1% of the raw milk samples analysed at concentrations between 0.025 and 1.01 µg/kg. Camaj et al.(2018), in Kosovo, detected AFM₁ in 38% of milk samples analysed. From 2013 to 2015, Zheng et al. (2017), in China, found out a mean of 21.2% of contaminated samples.

However, in Italy, several authors (Decastelli et al. 2007; Armorini et al. 2016; De Roma et al. 2017) reported very low levels of contamination in cow milk and usually no sample exceeded the EU regulation limit.

By contrast, a wide number of authors detected higher percentages of contaminated samples with concentrations that frequently exceeded the legal values. Bakirci (2001) in Turkey, found AFM₁ in 87.8% of the samples examined and with 44.3% of the cases surpassing the EU limit. In Ethiopia, Gizachew et al. (2016) reported that all the milk samples analysed were positive for the presence of AFM₁ and only 8.2% contained less than or equal to 0.05 µg/kg of AFM₁. Mohammed et al. (2016) in Tanzania detected that 83.8% of the milk samples were contaminated with AFM₁, exceeding all of them the EU established limit. These differences could be attributed to several factors including different analytical techniques, sample size and composition, season of the year, livestock management and dairy processing systems.

Carry-over of AFB₁ from feed to AFM₁ in milk

AFB₁ is transformed to AFM₁ in milk when lactating animals are fed with contaminated feed. However, only a small proportion of AFB₁ in feed leads to AFM₁ in the milk of dairy cows (Pittet 1998). Consequently, the incidence of AFB₁ in feed is much higher than the incidence of AFM₁ in raw milk.

Our study gave as a result that only fifteen raw milk samples positive for the presence of AFM₁ were obtained from cows that had been fed with contaminated feed having, consequently, a moderate positive correlation between AFM₁ contamination in milk and the level of AFB₁ contamination in feed. The carry-over rate varied from 0.6 to 6%, in agreement with data reported in other studies (Veldman et al. 1992; Pittet 1998; Unusan 2006; Iqbal et al. 2015). Other authors found lower rates of transformation that ranged from 1 to 3% (Diaz et al. 2004; van Eijkeren et al. 2006; Masoero et al. 2007). Sumantri et al. (2012) established that the carry-over was only 0.1%.

Despite all feed samples presented concentrations of AFB₁ below the EU limit, three raw milk samples exceeded the legal value for AFM₁. Moreover, 27 milk samples were positive for AFM₁ while correspondent feed samples were not contaminated with detectable levels of AFB₁. Conversely, the presence of AFB₁ in feed samples was not always reflected in milk. This fact has also been previously reported by other authors (Battacone et al., 2003; Gizachew et al., 2016; Han et al., 2013). Therefore, analysed levels of AFB₁ in feed within the limits do not guarantee a content of AFM₁ in milk under the permitted concentrations. A possible explanation could be the heterogeneous distribution of mycotoxins in raw materials and the problems derived from the sampling procedure. Only some sections of the feedstuffs are likely to be highly contaminated whereas most of the feed will probably not present any mycotoxin (Miraglia et al. 2005). Consequently, the amount of toxin could be underestimated if the contaminated fractions are not analysed. In a like manner, if only the heavily AF-contaminated fraction is examined, the mycotoxin content could be overestimated. Therefore, the sampling step may represent one of the main sources of variability in mycotoxin analyses. This procedure is carried out using a small proportion of the test sample (in our study 5 g from a bulk sample of 4-5 kg previously ground and homogenised) which is assumed to have the same mycotoxin concentration, but, even these subsamples could be different one from each other. Both, subsampling and grinding are also sources of error. The sampling error in toxin quantification has been shown to depend on the size of the sample. The error is reduced with bigger samples and subsamples and also with a smaller size of the ground particle (Whitaker 2003; Miraglia et al. 2005; Whitlow et al. 2010). Consequently, in order to try to minimise this error as much as possible, feed samples were dried and ground before being analysed and each sample was tested in triplicate.

TMR composition

Concentrates used in order to enrich feed for dairy cattle can represent up to a 70% of the daily feed ration, in our case they accounted for up to 40%. As a consequence, this ingredient may be an important source of mycotoxins due to its possible contamination with AFs. These toxins can be also found in cereal grains or

soybean products as well as in by-products from oil plants such as peanuts, sunflower seed, cotton seeds, palm kernels and copra (Nawaz et al. 1997; Scudamore and Livesey 1998; Placinta et al. 1999). Another source of AFs in the diet of dairy cows comes from the consumption of preserved feeding stuffs like silage, hay or straw (O'Brien et al. 2005; Mansfield and Kuldau 2007). After a long storage period, silage can be spoiled by some fungal species, being *Aspergillus* species (*A. fumigatus* and *A. flavus*) some of them (Cole et al. 1977). However, AFs only have been sporadically detected at low levels in forages. Maize has been traditionally related with the presence of AFs in dairy cattle diet, too (Whitlow and Hagler Jr. 2005), and it is usually one of the main components of animal feed (EFSA, 2013; Battilani et al., 2012).

The multivariate PLS regression carried out led us to identify which components of the total ration could have influenced on the presence of AFs in the feed (Figure 2 and Figure 3). Table 4 shows those raw materials that were positive for the presence of AFs and the range of concentrations. Positive samples of corn and compound feed could confirm the results of the PLS analysis in which it was observed that these ingredients seemed to be related with the presence of AFB₁ in the feed samples (Figure 2B and Figure 3A). The highest concentration of AFB₁ (2.44 µg/kg) was found in one of the compound feeds which was also positive for AFB₂, AFG₁ (3.25 µg/kg) and AFG₂. In this case, it was also analysed the correspondent TMR and it was found out to be positive for the presence of AFB₁, AFB₂ and AFG₁. In addition, milk samples from cows fed with this feed were contaminated with AFM₁. Some samples of cotton seed and corn were also positive for the presence of AFs. However, although cotton seed presented sometimes high concentrations of AFs, the toxins did not appear in the correspondent TMR sample. It may be explained because compound feed used to represent a relatively high proportion of the whole whereas the cotton seed represented normally a small portion (0.68-5.51%) of the total ration (Table 1).

Our positive outcomes in cotton seed and in corn are in agreement with other studies related to the analysis of feed ingredients. Liu et al. (2016) found that cottonseed was the most seriously contaminated feed ingredient examined. They also found that a high percentage of corn samples presented AFB₁. Sadegh et al. (2013) in Iran found out that cotton seed and sunflower meal samples were the most contaminated materials with AFB₁. The toxin was also presented in alfalfa, straw, rapeseed, cotton seed, corn silage and soybean meal samples although the highest concentration was in cotton seed. For Pleadin et al. (2015), maize was the most contaminated component although they also had positive AFB₁-samples of wheat, barley and oat. Whole cottonseed, as well as other oilseed such as soybean, flaxseed and sunflower, could be used as fat sources in dairy cattle feed. These ingredients allow a slow release of lipids in the rumen. Whole cottonseed has 22% crude protein, 20% ether

extract and 83% of total digestive nutrients, being, therefore, an excellent source of supplemental fat to increase diet energy density. Moreover, it seemed to be highly effective in maintaining milk fat. Despite all of these advantages, it could be related to the presence of AFs, therefore, it should be studied if it would be possible the replacement of **these** ingredient with others with similar nutritional properties (De Almeida et al. 2017).

Therefore, the level of contamination, both, in feed and milk was relatively low, in agreement with other studies carried out recently and this is probably related to the increasing surveillance measures and quality control of raw materials used for manufacturing feeds as well as the application of Good Agricultural and Storage Practices. However, in recent years, there are still outbreaks of AFM1 in milk over the legal limit and this may be linked to the raw materials used in the formulation of feedstuffs, **hence** it is important to have **present** information about the contribution of the ingredients on the AFB₁ content of the total mixed ration (TMR) given to cattle. Moreover, it is important to remember that, besides AFs, there are other mycotoxins that can occur in dairy feedstuffs and feed ingredients, including deoxynivalenol, zearalenone and fumonisins, that may be also harmful to animals and humans (Driehuis et al. 2008). Thus, mycotoxin contamination in feed and in milk should be regularly monitored, in particular the most susceptible ingredients, in order to reduce their incidence and their effects.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgments

This work was supported by the Spanish Ministry of Economy and Competitiveness (project AGL2014-55379-P). M. Rodríguez-Blanco thanks the University of Lleida for a pre-doctoral grant.

References

- Armorini S, Altafini A, Zaghini A, Roncada P (2016) Occurrence of aflatoxin M1 in conventional and organic milk offered for sale in Italy. *Mycotoxin Res* 32:237–246. doi: 10.1007/s12550-016-0256-8
- Bakirci I (2001) A study on the occurrence of aflatoxin M1 in milk and milk products produced in Van province of Turkey. *Food Control* 12:47–51. doi: 10.1016/S0956-7135(00)00020-7
- Bakirdere S, Bora S, Bakirdere EG, Aydin F, Arslan Y, Komesli OT, Aydin I, Yildirim E (2012) Aflatoxin species: Their health effects and determination methods in different foodstuffs. *Cent Eur J Chem* 10:675–

685. doi: 10.2478/s11532-012-0009-2
- Battacone G, Nudda A, Cannas A, Cappio Borlino A, Bomboi G, Pulina G (2003) Excretion of aflatoxin M1 in milk of dairy ewes treated with different doses of aflatoxin B1. *J Dairy Sci* 86:2667–2675. doi: 10.3168/jds.S0022-0302(03)73862-4
- Battilani P, Rossi V, Giorni P, Pietri A, Gualla A, Fels- Klerx HJ, Booij CJH, Moretti A, Logrieco A, Miglietta F, Toscano P, Miraglia M, Santis BDe, Brera C (2012) Modelling, predicting and mapping the emergence of aflatoxins in cereals in the EU due to climate change. *EFSA Support Publ* 9:1–172. doi: 10.2903/sp.efsa.2012.EN-223
- Camaj A, Meyer K, Berisha B, Arbneshti T, Haziri A (2018) Aflatoxin M1 contamination of raw cow's milk in five regions of Kosovo during 2016. *Mycotoxin Res* 34:205–209. doi: 10.1007/s12550-018-0315-4
- Cole RJ, Kirksey JW, Dorner JW, Wilson DM, Johnson Jr. J, Bedell D, Springer JP, Chexal KK, Clardy J, Cox RH (1977) Mycotoxins produced by *Aspergillus fumigatus* isolated from silage. *Ann la Nutr l'alimentation* 31:685–691
- De Almeida GF, Del Valle TA, De Paiva PG, De Jesus EF, Barletta RV, Gandra JR, Bettero VP, Takiya CS, Rennó FP (2017) Effects of whole raw soybean or whole cottonseed on milk yield and composition, digestibility, ruminal fermentation and blood metabolites of lactating dairy cows. *Anim Prod Sci* 57:122–128. doi: 10.1071/AN15266
- De Roma A, Rossini C, Ritieni A, Gallo P, Esposito M (2017) A survey on the Aflatoxin M1 occurrence and seasonal variation in buffalo and cow milk from Southern Italy. *Food Control* 81:30–33. doi: 10.1016/j.foodcont.2017.05.034
- Decastelli L, Lai J, Gramaglia M, Monaco A, Nachtmann C, Oldano F, Ruffier M, Sezian A, Bandirola C (2007) Aflatoxins occurrence in milk and feed in Northern Italy during 2004–2005. *Food Control* 18:1263–1266. doi: 10.1016/j.foodcont.2006.08.006
- Diaz DE, Hagler Jr. WM, Blackwelder JT, Eve JA, Hopkins BA, Anderson KL, Jones FT, Whitlow LW (2004) Aflatoxin Binders II: Reduction of aflatoxin M1 in milk by sequestering agents of cows consuming aflatoxin in feed. *Mycopathologia* 157:233–241. doi: 10.1023/B:MYCO.0000020587.93872.59
- Driehuis F, Spanjer MC, Scholten JM, te Giffel MC (2008) Occurrence of Mycotoxins in Feedstuffs of Dairy

- Cows and Estimation of Total Dietary Intakes. *J Dairy Sci* 91:4261–4271. doi: 10.3168/jds.2008-1093
- El Mundo (2013). In: El Mundo.
- <http://www.elmundo.es/andalucia/2013/11/28/52971de963fd3dc5318b458d.html>. Accessed 11 Jan 2019
- European Commission (EC) (2002) Commission Directive 2002/32/EC of 7 May 2002 on undesirable substances in animal feed. *Off J Eur Union* L140:10–21
- European Commission (EC) (2006) Commission Regulation 1881/2006 of 19 December 2006 setting maximum levels for certain contaminants in foodstuffs. *Off J Eur Union* L364:5–24
- European Food Safety Authority (EFSA) (2013) Aflatoxins (sum of B1, B2, G1, G2) in cereals and cereal-derived food products. *EFSA Support Publ* 10:1–11. doi: 10.2903/sp.efsa.2013.EN-406
- Ehsani A, Barani A, Nasiri Z (2016) Occurrence of aflatoxin B1 contamination in dairy cows feed in Iran. *Toxin Rev* 35:54–57. doi: 10.3109/15569543.2016.1155622
- Fallah AA, Jafari T, Fallah A, Rahnama M (2009) Determination of aflatoxin M1 levels in Iranian white and cream cheese. *Food Chem Toxicol* 47:1872–1875. doi: 10.1016/j.fct.2009.04.042
- Food and Agriculture Organization (FAO) (2004) Worldwide regulations for mycotoxins in food and in feed in 2003. Rome (Italy)
- Food and Drug Administration (FDA) (1996) Action levels for aflatoxins in animal feeds, Compliance Policy Guides CPG Sec 683.100.
- <https://www.fda.gov/iceci/compliancemanuals/compliancepolicyguidancemanual/ucm074703.htm>. Accessed 11 Jan 2019
- Fink-Gremmels J (1999) Mycotoxins: Their implications for human and animal health. *Vet Q* 21:115–120. doi: 10.1080/01652176.1999.9695005
- Garrido CE, Hernández Pezzani C, Pacin A (2012) Mycotoxins occurrence in Argentina's maize (*Zea mays* L.), from 1999 to 2010. *Food Control* 25:660–665. doi: 10.1016/j.foodcont.2011.11.043
- Gizachew D, Szonyi B, Tegegne A, Hanson J, Grace D (2016) Aflatoxin contamination of milk and dairy feeds in the Greater Addis Ababa milk shed, Ethiopia. *Food Control* 59:773–779. doi: 10.1016/j.foodcont.2015.06.060

- Golge O (2014) A survey on the occurrence of aflatoxin M1 in raw milk produced in Adana province of Turkey. Food Control 45:150–155. doi: 10.1016/j.foodcont.2014.04.039
- Han RW, Zheng N, Wang JQ, Zhen YP, Xu XM, Li SL (2013) Survey of aflatoxin in dairy cow feed and raw milk in China. Food Control 34:35–39. doi: 10.1016/j.foodcont.2013.04.008
- Hernández-Martínez R, Navarro-Blasco I (2015) Surveillance of aflatoxin content in dairy cow feedstuff from Navarra (Spain). Anim Feed Sci Technol 200:35–46. doi: 10.1016/j.anifeedsci.2014.12.002
- Heshmati A, Milani JM (2010) Contamination of UHT milk by aflatoxin M1 in Iran. Food Control 21:19–22. doi: 10.1016/j.foodcont.2009.03.013
- Horwitz W, Latimer GW (2006) Official methods of analysis of AOAC International, 18 th edit. Gaithersburg, Maryland
- International Agency for Research on Cancer (IARC) (2012) Monograph on the evaluation of carcinogenic risk to humans: Chemical agents and related occupations. A review of humans carcinogens. Lyon (France)
- International Agency for Research on Cancer (IARC) (1993) Some naturally occurring substances: food items and constituents, heterocyclic aromatic amines and mycotoxins. Lyon (France)
- Iqbal SZ, Asi MR, Jinap S (2013) Variation of aflatoxin M1 contamination in milk and milk products collected during winter and summer seasons. Food Control 34:714–718. doi: 10.1016/j.foodcont.2013.06.009
- Iqbal SZ, Jinap S, Pirouz AA, Ahmad Faizal AR (2015) Aflatoxin M1 in milk and dairy products, occurrence and recent challenges: A review. Trends Food Sci. Technol. 46:110–119
- Ismail A, Riaz M, Akhtar S, Yoo SH, Park S, Abid M, Aziz, M, Ahmad, Z (2017) Seasonal variation of aflatoxin B1 content in dairy feed. J Anim Feed Sci 26:33–37. doi: 10.22358/jafs/69008/2017
- International Standard Organization (ISO) (2007) ISO 14501:2007 (IDF 171:2007). Milk and milk powder -- Determination of aflatoxin M1 content -- Clean-up by immunoaffinity chromatography and determination by high-performance liquid chromatography.
- Juan C, Soriano JM, Burdaspal P (2007) Aflatoxinas del grupo B y G. In: Díaz de Santos (ed) Micotoxinas en alimentos. Madrid (Spain), pp 167–184
- Kocasari FS, Tasci F, Mor F (2012) Survey of aflatoxin M1 in milk and dairy products consumed in Burdur,

- Turkey. *Int J Dairy Technol* 65:365–371. doi: 10.1111/j.1471-0307.2012.00841.x
- Liu J, Sun L, Zhang J, Guo J, Chen L, Qi D, Zhang N (2016) Aflatoxin B1, zearalenone and deoxynivalenol in feed ingredients and complete feed from central China. *Food Addit Contam Part B Surveill* 9:91–97. doi: 10.1080/19393210.2016.1139003
- Mansfield MA, Kuldau GA (2007) Microbiological and molecular determination of mycobiota in fresh and ensiled maize silage. *Mycologia* 99:269–278. doi: 10.3852/mycologia.99.2.269
- Martins HM, Guerra MMM, Bernardo FMD (2007) Occurrence of aflatoxin B in dairy cow's feed over 10 years in Portugal (1995-2004). *Rev Iberoam Micol* 24:69–71
- Martins ML, Martins HM (2000) Aflatoxin M1 in raw and ultra high temperature-treated milk commercialized in Portugal. *Food Addit Contam* 17:871–874
- Masoero F, Gallo A, Moschini M, Piva G, Diaz D (2007) Carryover of aflatoxin from feed to milk in dairy cows with low or high somatic cell counts. *Animal* 1:1344–1350. doi: 10.1017/S1751731107000663
- Miraglia M, De Santis B, Minardi V, Debegnach F, Brera C (2005) The role of sampling in mycotoxin contamination: An holistic view. *Food Addit Contam* 22:31–36. doi: 10.1080/02652030500389055
- Mohammed S, Munissi JJE, Nyandoro SS (2016) Aflatoxin M1 in raw milk and aflatoxin B1 in feed from household cows in Singida, Tanzania. *Food Addit Contam Part B Surveill* 9:85–90. doi: 10.1080/19393210.2015.1137361
- Nawaz S, Scudamore KA, Rainbird SC (1997) Mycotoxins in ingredients of animal feeding stuffs: I. Determination of *Alternaria* mycotoxins in oilseed rape meal and sunflower seed meal. *Food Addit Contam* 14:249–262
- O'Brien M, O'Kiely P, Forristal PD, Fuller HT (2005) Fungi isolated from contaminated baled grass silage on farms in the Irish Midlands. *FEMS Microbiol Lett* 247:131–135. doi: 10.1016/j.femsle.2005.04.037
- Pittet A (1998) Natural occurrence of mycotoxins in foods and feeds - An updated review. *Rev Med Vet (Toulouse)* 149:479–492
- Placinta CM, D'Mello JPF, MacDonald AMC (1999) A review of worldwide contamination of cereal grains and animal feed with *Fusarium* mycotoxins. *Anim Feed Sci Technol* 78:21–37. doi: 10.1016/S0377-8401(98)00278-8

- Pleadin J, Vulić A, Perši N, Škrivanko M, Capek B, Cvetnić T (2015) Annual and regional variations of aflatoxin B1 levels seen in grains and feed coming from Croatian dairy farms over a 5-year period. *Food Control* 47:221–225. doi: 10.1016/j.foodcont.2014.07.017
- Sadegh M, Sani AM, Ghiasvand R (2013) Determination of aflatoxin B1 in animal feed in Mashhad, Iran. *Biotechnol An Indian J* 7:334–336
- Sahin HZ, Celik M, Kotay S, Kabak B (2016) Aflatoxins in dairy cow feed, raw milk and milk products from Turkey. *Food Addit Contam Part B Surveill* 9:152–158. doi: 10.1080/19393210.2016.1152599
- Sassahara M, Pontes Netto D, Yanaka EK (2005) Aflatoxin occurrence in foodstuff supplied to dairy cattle and aflatoxin M1 in raw milk in the North of Paraná state. *Food Chem Toxicol* 43:981–984. doi: <https://doi.org/10.1016/j.fct.2005.02.003>
- Scudamore KA, Livesey CT (1998) Occurrence and significance of mycotoxins in forage crops and silage: A review. *J Sci Food Agric* 77:1–17. doi: 10.1002/(SICI)1097-0010(199805)77:1<1::AID-JSFA9>3.0.CO;2-4
- Sumantri I, Murti TW, Van Der Poel AFB, Boehm J, Agus A (2012) Carry-over of aflatoxin B1-feed into aflatoxin M1-milk in dairy cows treated with natural sources of aflatoxin and bentonite. *J Indones Trop Anim Agric* 37:271–277
- Unusan N (2006) Occurrence of aflatoxin M1 in UHT milk in Turkey. *Food Chem Toxicol* 44:1897–1900. doi: 10.1016/j.fct.2006.06.010
- van Eijkeren JCH, Bakker MI, Zeilmaker MJ (2006) A simple steady-state model for carry-over of aflatoxins from feed to cow's milk. *Food Addit Contam* 23:833–838. doi: 10.1080/02652030600779890
- Veldman A, Meijs JAC, Borggreve GJ, Heeres-Van Der Tol JJ (1992) Carry-over of aflatoxin from cows' food to milk. *Anim Prod* 55:163–168. doi: 10.1017/S0003356100037417
- Whitaker TB (2003) Detecting mycotoxins in agricultural commodities. *Mol Biotechnol* 23:61–71. doi: 10.1385/MB:23:1:61
- Whitlow LW, Hagler Jr. WM, Díaz DE (2010) Mycotoxins in feeds. In: *Feedstuffs*. <https://pdfs.semanticscholar.org/66a9/4da4a8fb260176cfd1fab196001a30851f1.pdf>. Accessed 11 Jan 2019

Whitlow LW, Hagler Jr. WM (2005) Mycotoxins in Dairy Cattle: Occurrence, Toxicity, Prevention and Treatment. In: Proc. Southwest Nutr. Conf.

<http://www.dairyweb.ca/Resources/USWebDocs/Mycotoxins4.pdf>. Accessed 11 Jan 2019

Zheng N, Li SL, Zhang H, Min L, Gao YN, Wang JQ (2017) A survey of aflatoxin M1 of raw cow milk in China during the four seasons from 2013 to 2015. Food Control 78:176–182. doi: 10.1016/J.FOODCONT.2017.02.055

Figure captions

Fig. 1 AFM₁-positive milk samples from February 2016 to January 2018

Fig. 2 Importance of the TMR components (VIP) on the presence of AFs (A), of AFB₁ (B) and of AFG₁ (C) in feed samples

Fig. 3 Influence of the TMR components (VIP) on AFB₁ presence in feed (A) and on AFM₁ in milk from cows fed with feed (B) taking into account the ingredients of the compound feed

Tables

Table 1. Main ingredients of total mixed ration samples, number of samples that contained each of them and percentage of inclusion in the feed.

TMR component	No. of samples (% total)	Percentage (%)
Compound feed	145 (75.13%)	1.61-59.18
Maize silage	107 (55.44%)	9.92-76.92
Straw	95 (49.22%)	0.68-10.71
Dehydrated alfalfa	78 (40.41%)	2.28-25.66
Alfalfa hay	63 (32.64%)	1.63-32.65
Corn	51 (26.42%)	0.11-22.83
Grass silage	47 (24.35%)	9.89-89.29
Bagasse	40 (20.73%)	3.56-25.81
Immature corn silage	33 (17.10%)	3.76-19.05
Sugar beet pulp	30 (15.54%)	0.93-22.79
Cotton seed	30 (15.54%)	0.68-5.51
Ray grass silage	28 (14.51%)	7.60-75.95
Alfalfa silage	21 (10.88%)	5.71-44.78
Soya bean	21 (10.88%)	3.04-14.47
Barley	21 (10.88%)	1.13-8.10

Vetch hay	15 (7.77%)	2.06-13.64
Oat silage	13 (6.74%)	2.00-47.46
Dehydrated ray-grass	12 (6.12%)	1.90-8.44
Soya bean husk	10 (5.18%)	1.49-2.54
Barley silage	8 (4.14%)	16.01-23.25
Wheat silage	5 (2.59%)	18.60-32.38
Okara	4 (2.07%)	3.72-8.85

Table 2. Aflatoxins in feed samples (n=193).

Year	Analyte	No. of samples	No. of positive samples (%)	Range (µg/kg)
2016	AFB ₁	103	16 (15.53%)	0.14-4.66
	AFB ₂		5 (4.85%)	0.05-0.41
	AFG ₁		8 (7.77%)	0.14-1.35
	AFG ₂		7 (6.80%)	0.1-0.57
2017	AFB ₁	79	7 (8.86%)	0.14-1.82
	AFB ₂		3 (3.80%)	0.08-0.20
	AFG ₁		32 (40.51%)	0.14-6.45
	AFG ₂		6 (7.59%)	0.1-0.47
2018	AFB ₁	11	1 (9.10%)	0.27
	AFB ₂		1 (9.10%)	0.06
	AFG ₁		7 (63.64%)	0.13-0.45
	AFG ₂		0	-

Table 3. Aflatoxin M₁ in milk samples obtained 24 and 48 h after feed consumption (n=375).

Year	Analyte	No. of samples	No. of positive samples (%)	Range (µg/kg)
2016	AFM ₁ (24 h)	102	30 (29.41%)	0.009-1.36
	AFM ₁ (48 h)	102	28 (27.45%)	0.009-0.648
2017	AFM ₁ (24 h)	74	4 (5.41%)	0.015-0.024
	AFM ₁ (48 h)	78	7 (8.97%)	0.010-0.032
2018	AFM ₁ (24 h)	8	1 (12.5%)	0.009
	AFM ₁ (48 h)	11	1 (9.09%)	0.011

Table 4. Aflatoxin-positive raw materials analysed separately.

Raw material	No. of total samples	No. of positive samples (range, µg/kg)			
		AFB ₁	AFB ₂	AFG ₁	AFG ₂
Ray grass silage	1	0	0	0	1 (0.26)
Corn	6	3 (0.24-1.05)	1 (0.06)	3 (0.18-0.76)	2 (0.11-0.78)
Cotton seed	12	5 (0.08-1.97)	2 (0.16-0.21)	4 (0.27-3.68)	3 (0.23-0.77)
Soya bean	2	1 (0.26)	0	0	0
Compound feed	5	2 (0.17-2.44)	2 (0.05-0.21)	3 (0.11-3.25)	2 (0.16-0.2)

Figure 1

Figure 2

Figure 3