Universitat de Lleida

Document downloaded from:

<http://hdl.handle.net/10459.1/69990>

The final publication is available at:

<https://doi.org/10.1111/nph.16656>

Copyright

(c) Patsiou, Theofania et al., 2020

(c) New Phytologist Trust, 2020

New Phytologist

Intraspecific responses to climate reveal nonintuitive warming impacts on a widespread thermophilic conifer

Journal:	<i>New Phytologist</i>
Manuscript ID	NPH-MS-2019-32006.R2
Manuscript Type:	MS - Regular Manuscript
Date Submitted by the Author:	28-Apr-2020
Complete List of Authors:	Patsiou, Theofania-Sotiria; University of Basel, Umweltwissenschaften Shestakova, Tatiana; Woods Hole Research Center, Department of Plant and Environmental sciences Klein, Tamir; Weizmann Institute of Science, Department of Plant and Environmental Sciences di Matteo, Giovanni; CREA, Council for Agricultural Research and Economics Bay, Hassan; Forest Research center, (CRF) Chambel, Maria; Instituto Nacional de Investigacion y Tecnologia Agraria y Alimentaria, Ecologia y genetica forestal Zas, Rafael; Mision Biologica de Galicia, Superior de Investigaciones Científicas Voltas, Jordi; University of Lleida, Department of Crop and Forest Science
Key Words:	Aleppo pine, climate sensitivity, ecotype, factorial regression, intraspecific adaptation, response function, space for time substitution

SCHOLARONE™
Manuscripts

Type of paper: Primary Research Article

Intraspecific responses to climate reveal nonintuitive warming impacts on a widespread thermophilic conifer

Running Head: Ecotypic responses of a thermophilic conifer

Theofania S. Patsiou¹, Tatiana A. Shestakova², Tamir Klein³, Giovanni di Matteo⁴, Hassan Sbaya⁵, M^a Regina Chambel⁶, Rafael Zas⁷, Jordi Voltas^{8,9}

¹Institute of Botany, University of Basel, Schönbeinstrasse 6, CH-4056 Basel, Switzerland

²The Woods Hole Research Center, 149 Woods Hole Road, Falmouth, MA 02540, USA

³Department of Plant and Environmental sciences, Weizmann Institute of Science, 234 Herzl Street, Rehovot 7610001, Israel

⁴Council for Agricultural Research and Economics, Research Centre for Agriculture and Environment (CREA), via della Navicella 2-4, 00186, Rome, Italy

⁵Forest Research Centre (CRF), Av. Omar Ibn el Khattab. Agdal, 110000 Rabat, Morocco

⁶CIFOR-INIA, Ctra.de La Coruña km 7.5, E-28040 Madrid, Spain

⁷Misión Biológica de Galicia, Consejo Superior de Investigaciones Científicas (MBG-CSIC), Apdo. 28, E-36080 Salcedo, Spain

⁸Joint Research Unit CTFC - AGROTECNIO, Av. Alcalde Rovira Roure 191, E-25198 Lleida, Spain

⁹Department of Crop and Forest Sciences, University of Lleida, Av. Alcalde Rovira Roure 191, E-25198 Lleida, Spain

Corresponding author: Jordi Voltas, jvoltas@pvcf.udl.cat

ORCID ID 0000-0003-4051-1158

Total word count for the main body of the text: 6405

Word counts for each section: Introduction: 1211; Materials and Methods: 1760; Results: 1610; Discussion: 1824

Number of Figs. (5); Tables: (1); Supporting Information: Figs. (7); Tables (12).

Summary

- Many ecologically important forest trees from dry areas have been insufficiently investigated for their ability to adapt to the challenges posed by climate change, which hampers the implementation of mitigation policies. We analyzed 14 common-garden experiments across the Mediterranean of the widespread thermophilic conifer *Pinus halepensis*, involving 157 populations categorized into five ecotypes.
- Ecotype-specific tree height responses to climate were applied to projected climate change (2071-2100 AD), to project potential growth patterns both locally and across the species' range.
- We found contrasting ecotypic sensitivities to annual precipitation but comparatively uniform responses to mean temperature, while evidence of local adaptation for tree height was limited to mesic ecotypes. We projected intriguing patterns of response range-wide, implying either height inhibition or stimulation of up to 75%, and deduced that the ecotype currently experiencing more favorable (wetter) conditions will show the largest inhibition. Extensive height reductions can be expected for coastal areas of France, Greece, Spain and northern Africa.
- Our findings underline that intraspecific variations in sensitivity to precipitation must be considered when projecting tree height responses of dry forests to future climate. The ecotype-specific projected performances call for management activities to ensure forest resilience in the Mediterranean through, e.g. tailored deployment strategies.

Key words: Aleppo pine, climate sensitivity, ecotype, factorial regression, intraspecific adaptation, response function, space for time substitution

66 Introduction

67 Trees are crucial organisms in many terrestrial ecosystems, and their responses to
 68 climate variations could influence the global carbon and water cycles considerably and
 69 affect the ecosystem services that forests provide (Viglizzo *et al.*, 2016; Sheil, 2018).
 70 Conifers, in particular, show strongly differentiated populations (Neale & Wheeler,
 71 2019) that result from long-term selective processes, the history of niche recolonization
 72 after glaciations, and the trade-offs derived from patterns of covariation among fitness-
 73 related traits (Kawecki & Ebert, 2004). High levels of intraspecific genetic variability
 74 are essential for the resilience of forests (Alberto *et al.*, 2013). However, the importance
 75 of climatic adaptation of forest trees, which involves both standing genetic variation and
 76 differential phenotypic plasticity of populations to buffer environmental changes
 77 (Mátyás, 1996), has been largely overlooked when forecasting responses to climate
 78 change. This omission may result in unrealistic projections for the status and
 79 distribution of forest species in the forthcoming decades (O'Neill *et al.*, 2008; Reich &
 80 Oleksyn, 2008; Peterson *et al.*, 2019).

81 The analysis of adaptive strategies has focused primarily on forest trees of high
 82 economic relevance in the context of breeding programs (e.g., *Pinus contorta*; Montwé
 83 *et al.*, 2016). However, the ability of many ecologically important species to adapt to
 84 the challenges posed by climate change remains insufficiently studied (Mátyás *et al.*,
 85 2009). This lack of knowledge precludes the implementation of policies and measures
 86 for climate change mitigation and adaptation. Besides, these species tend to occupy
 87 peripheral, arid areas that are subject to severe human impact. One such example is the
 88 circum-Mediterranean conifer Aleppo pine (*Pinus halepensis* Mill.). This thermophilic
 89 pine is the most widely distributed tree species in the Mediterranean basin. It shows an
 90 extensive ecological breadth and is seemingly adapted to a broad range of abiotic
 91 stressors and disturbances (Ne'eman *et al.*, 2004; Schiller & Atzmon, 2009; Klein *et al.*,
 92 2013; Santini *et al.*, 2019). Previous studies have shown that populations in xeric sites
 93 exhibit different characteristics than those in mesic areas, indicating intricate
 94 anatomical, morphological, and physiological adaptations of the species (Santos del
 95 Blanco *et al.*, 2013; Voltas *et al.*, 2015; David-Schwartz *et al.*, 2016). However, a
 96 comprehensive analysis of the climate factors underlying intraspecific variation is
 97 currently lacking for *P. halepensis*, which exemplifies the significant gaps in our
 98 knowledge on the adaptive patterns of tree species and populations inhabiting drought-
 99 prone regions (Mátyás *et al.*, 2009).

Local adaptation, which reflects a genetic adjustment of individuals to a particular environmental niche (i.e., higher fitness of resident than of foreign genotypes; Williams [1966]), is widespread in forest trees (Alberto *et al.*, 2013; Valladares *et al.*, 2014; Peterson *et al.*, 2019; Sáenz-Romero *et al.*, 2019). The first step towards detecting local adaptation is to find evidence that a trait is relevant for fitness, which can be suggested by linking trait variability of populations with the historical climate at their location of origin. This has been extensively shown for tree height and diameter in the case of *P. halepensis* (Climent *et al.*, 2008; Voltas *et al.*, 2008; Esteban *et al.*, 2010), which suggests ecotypic specialization for either resource-rich habitats favoring vegetative growth, or resource-poor habitats favoring early reproduction. However, local adaptation is challenging to characterize because of the differential allocation of resources to reproduction, storage, defence, and above- and below-ground growth throughout ontogeny trades-off for maximizing fitness, as evidenced for *P. halepensis* (Sbay & Zas, 2018; Santini *et al.*, 2019). Despite these difficulties, height growth is the most widely used fitness-related trait to date for understanding adaptation in forest trees (Fréjaville *et al.*, 2020; Peterson *et al.*, 2019), mainly because of its ample availability in extensive networks of common gardens. In particular, tree height can be considered a potential proxy for fitness in *P. halepensis* because this species is shade-intolerant, thus showing light-dependent growth patterns, and presents phenotypic plasticity subject to resource availability (Zavala *et al.*, 2011).

The analysis and interpretation of intraspecific adaptive patterns can be accomplished using common-garden experiments and thorough statistical modeling of genotype by environment (G×E) interactions (Mátyás, 1996; Rehfeldt *et al.*, 2002). Within this context, the analysis of climate-response-functions, which describe the relationships between the performance of a population across test sites and the climate of those sites (Rehfeldt *et al.*, 2002), can be useful in forecasting the effects of climate change on tree species (Peterson *et al.*, 2019). Thus, the response of populations to the new environments at the test sites can be interpreted as a simulation of their response to climate change. In this study, climate-response-functions were obtained using a mixed factorial regression model approach. This approach is often used in plant breeding given the relevance of G×E for the optimal deployment of new releases (van Eeuwijk *et al.*, 2005; Malosetti *et al.*, 2013; Heslot *et al.*, 2014), but is less frequently applied in forest genetics (Sixto *et al.*, 2016). Factorial regression describes G×E interactions as differential genotypic sensitivities to specific environmental factors (i.e., differences in

the slope of response). This methodology has two main advantages in gaining insight into the environmental causes underlying adaptation. First, it allows for straightforward statistical testing of the drivers and relevance of differential plasticity linked to sizeable G×E interactions. Secondly, it is flexible in accounting for population structure (i.e., ecotypic variation) in a multi-environment setting, which is valuable for highly unbalanced datasets denoting incomplete replication of populations across testing sites.

We hypothesized that the combined responses to temperature and precipitation had shaped the intraspecific adaptive strategies of the widespread thermophilic pine, *P. halepensis*. We used tree height to unravel the potential of populations to perform across a range-wide environmental gradient. Out of the components of above-ground growth, tree height is under the strongest genetic control and, as such, is commonly used as an indicator of population and site differences in productivity (Skovsgaard & Vancley, 2007; Moles *et al.*, 2009). Our goal was to delineate ecotypic responses to climate and obtain projections of future growth under climate change, which are related to the carbon storage capacity of dry forests and their role as net carbon sinks (Zhu *et al.*, 2018). Under climate fluctuations, the productivity of species should decrease from recently favorable habitats, the 'leading-edge' where populations are expanding into, to the 'trailing-edge', where populations are contracting from recently unsuitable habitats, following a climate gradient.

Specifically, this study addresses the following questions: (i) what is the relative importance of precipitation and temperature as factors that could explain adaptive variation in a thermophilic species subject to varying levels of water shortage?, (ii) does accounting for G×E interaction improve the estimation accuracy of tree height for the current climate and modify the projections of tree height under climate change?, and (iii) are ecotypes from dry environments more likely to decline in growth by climate warming than those occupying wetter areas, hence following the commonly assumed 'trailing-edge, leading-edge' range-shift pattern? To answer these questions, we analyzed a broad, yet, highly unbalanced trial network in which five major *P. halepensis* ecotypes were evaluated across the Mediterranean, including independent data validation (new population-trial combinations). Our study is likely to be the first range-wide one examining adaptive variation for an ecologically relevant forest species native to warm, dry environments which uses independent data to validate predictions. Our study further analyzes the observed ecotypic patterns to forecast the species' response to future conditions (period 2071-2100 AD).

Materials and Methods

Field trials and plant material

Nine Aleppo pine provenance trials (hereafter, training trials) were used to obtain intraspecific response functions to climate. The training trials were distributed across the Mediterranean basin (Israel, Italy, and Spain; Fig. S1 in Supporting Information, SI), in areas representative of the climatic conditions where Aleppo pine is found naturally. In particular, MAT (mean annual temperature) ranged from 11.1 to 19.5 °C and MAP (mean annual precipitation) from 291 to 757 mm (high-resolution WorldClim database v. 1.4; Hijmans *et al.*, 2005). The climate of the training trials (SI Table S1) closely matched the species' core climate envelope, with an average difference between midpoint of training trials and the midpoint of the species distribution range of +1.3 °C (MAT) and -55 mm (MAP). The training trials (SI Table S2) had been previously used to characterize the extent of ecotypic differentiation in tree height plasticity and survival for the species, with no significant ecotype by trial interactions for height (Voltas *et al.*, 2018). Five additional trials with climate characteristics within the range of those of the training dataset were used as test trials to evaluate the predictive ability of response functions (see SI Tables S1 and S2) and were from Australia (Spencer, 1985), Italy, Morocco (Sbay & Zas, 2018) and Spain (Santos del Blanco *et al.*, 2013).

The training trials evaluated seed sources from 82 populations, representing most of the species' natural distribution across the Mediterranean basin. The test trials evaluated seed sources from 142 populations, with 67 populations being also present in the training trials (SI Fig. S1 and Table S3). Both datasets were extremely unbalanced (SI Materials and Methods, section A). A total of 157 different populations were considered. For each population, climatic variables at their origin were obtained for the period 1961–1990 (Climatic Research Unit, CRU TS 3.24; Harris *et al.*, 2014), at a coarse resolution (0.5°×0.5° grid-box basis) following Voltas *et al.* (2018), and were retrieved from each population's nearest grid. The climate variables were selected based on previous identification of climate factors underlying ecotypic variation (Tapias *et al.*, 2004; Voltas *et al.*, 2008). These included mean annual temperature (MAT_o), the maximum temperature of the warmest month (i.e., July; TMX_o), minimum temperature of the coldest month (i.e., January; TMN_o), the temperature annual range (i.e., TMX_o - TMN_o; TAR_o), mean annual precipitation (MAP_o), and mean summer precipitation (MSP_o) at the origin.

Grouping of training populations and assignment of test populations to ecotypes

Using hierarchical cluster analysis (Ward, 1963), we identified ecotypes developing in contrasting climates and, thus, putatively diverging via adaptations to heterogeneous conditions (Voltas *et al.*, 2018). Each population was assigned to a different ecotype based on climatic information (SI Table S3). Climate variables used were MAT_o, MAP_o, TAR_o, and MSP_o. The inclusion of two additional variables (TMX_o and TMN_o) did not substantially modify the classification (results not shown). The resulting five ecotypes (SI Fig. S2) were assigned to a particular group (climate type) based on three criteria, following (Le Hourou, 2004): (i) summer precipitation (dry summer: MSP_o ≤ 60 mm; wet summer: MSP_o ≥ 80 mm), (ii) annual precipitation (arid-semiarid: MAP_o ≤ 600 mm; sub-humid: MAP_o > 600), and (iii) winter temperature (cold: −1 < TMN_o ≤ 1C; cool: 1 < TMN_o ≤ 3C; temperate: 3 < TMN_o ≤ 5C). The groups were, therefore, identified as (a) Dry-summer/Semiarid/Temperate (DST, 23 populations), (b) Dry-summer/Semiarid/Cold (DSC, 24 populations), (c) Dry-summer/sub-Humid/Temperate (DHT, 12 populations), (d) Wet-summer/Semiarid/Temperate (WST, 18 populations), and (f) Wet-summer/sub-Humid/Cool (WHC, 5 populations); see SI Table S4 for the main climatic characteristics of ecotypes. Ecotypes from sub-humid climates (DHT, WHC) are, hereafter, referred to as mesic ecotypes (SI Table S5).

Using MAT_o, MAP_o, TAR_o and MSP_o, we obtained linear discriminant functions of the five ecotypes identified through clustering. The discriminant functions were employed to allocate each of the 75 new populations of the test dataset to a particular ecotype and to obtain subsequent probabilities of membership (SI Table S6).

Multi-environment trial (MET) analysis and factorial regression modeling

The nine training trials were combined and analyzed as multi-environment mixed model analysis of variance for population-trial means with the fixed trial (*T*), ecotype (*E*) and trial by ecotype interaction effects, and random effects for the population (*P*) nested to ecotype and population nested to ecotype by trial interaction (random terms are underlined):

$$Y_{ijk} = \mu + T_i + E_k + \underline{P(E)_{jk}} + (TE)_{ik} + \underline{TP(E)_{ijk}} \quad [1]$$

where Y_{ijk} is the observation (tree height) of the j^{th} population of the k^{th} ecotype in the i^{th} trial, μ is the general mean, T_i is the fixed effect of the i^{th} trial, E_k is the fixed effect of

the k^{th} ecotype, $P(E)_{jk}$ is the random effect of the j^{th} population nested to the k^{th} ecotype, $(TE)_{ik}$ is the fixed effect of interaction between the k^{th} ecotype and the i^{th} trial, and $TP(E)_{ijk}$ is the random (residual) effect of interaction between the j^{th} population nested to the k^{th} ecotype and the i^{th} trial.

For the explanation of ecotype by trial effects, we used a factorial regression modeling approach by which regression terms for the interpretation of genotype by environment (G×E) interaction are incorporated in the form of environmental covariates to the levels of the environmental factor. Factorial regression helps to detect genotypes that are differentially sensitive to changes in identified environmental components (van Eeuwijk, 1995; Malosetti *et al.*, 2013).

We fitted mixed factorial regression models (Denis *et al.*, 1997) to ecotype by trial effects in model [1] searching for climate variables underlying differential plastic responses at the ecotype level (Voltas *et al.*, 2018). In factorial regression, an explicit environmental covariable Z can be included in the levels of the trial factor to describe the trial main effect plus the interaction term of interest, $(TE)_{ik}$:

$$Y_{ijk} = \mu + \alpha Z_i + T_i + E_k + \underline{P(E)_{jk}} + \beta_k Z_i + (TE)_{ik} + \underline{TP(E)_{ijk}} \quad [2]$$

where α stands for the general sensitivity of all populations to the environmental variable Z_i , β_k stands for the specific sensitivity of the k^{th} ecotype to the same variable, and T_i and $(TE)_{ik}$ are the residual effects of the i^{th} trial and the interaction between the k^{th} ecotype and the i^{th} trial, respectively. Model [2] allows identification of a climate variable that can explain absolute changes in tree height among trials, differential ecotypic reactions across trials, or both simultaneously. This model can be extended to the multiple covariables case:

$$Y_{ijk} = \mu + \sum_{l=1}^L \alpha_l Z_{il} + T_i + E_k + \underline{P(E)_{jk}} + \sum_{l=1}^L \beta_{kl} Z_{il} + (TE)_{ik} + \underline{TP(E)_{ijk}} \quad [3]$$

where Z_{il} denotes the values for l climate variables underlying variation in T_i , $(TE)_{ik}$, or both. The terms $\alpha_l Z_{il}$ and $\beta_{kl} Z_{il}$ are fixed terms because Z_{il} values contain known, explicit environmental information.

Characterization of ecotypic responses to climate: response surfaces

We tested, for the selection of climatic factors in model [3], all possible combinations of one temperature- and one precipitation-related variable because temperature (or precipitation) variables were significantly correlated, hence, partly redundant for explaining differential ecotypic performances: MAT and MAP, MAT and MSP, TAR and MAP, and TAR and MSP. For each combination, a quadratic response surface was incorporated into the levels of the trial main effect and the ecotype by trial interaction simultaneously in [3], as follows:

$$\sum_{l=1}^2 \alpha_l Z_{il} = \alpha_1 Z_{i1} + \alpha_2 Z_{i2} + \alpha'_1 Z_{i1}^2 + \alpha'_2 Z_{i2}^2 + \alpha_{12} Z_{i1} Z_{i2} \quad [4]$$

and

$$\sum_{l=1}^2 \beta_{kl} Z_{il} = \beta_{k1} Z_{i1} + \beta_{k2} Z_{i2} + \beta'_{k1} Z_{i1}^2 + \beta'_{k2} Z_{i2}^2 + \beta_{k12} Z_{i1} Z_{i2} \quad [5]$$

Where Z_{il} denotes the values for $l=2$ climate variables of temperature and precipitation at the trial i . We refer to this model as a factorial regression response surface model (hereafter FRSM). The heterogeneity in the β_k 's for Z_l and Z_2 accounts for the interaction between ecotypes and trials, while the sum of multiplicative terms $\sum_{l=1}^2 \beta_{kl} Z_{il}$ approximates this interaction (van Eeuwijk, 1995). To facilitate interpretation of the β_k 's, the climate variables were standardized to mean zero and standard deviation of one. For the selection of the best combination of climate variables in the FRSM, we used the marginal R^2 (Nakagawa & Schielzeth, 2013), which informed on the variance explained by the fixed (climate) factors alone, i.e., the predictive ability of each model based on the (non-standardized) climate variables, and omitting any trial-related effect from model [3]. The data met the underlying assumptions of normality and homogeneity of error variances (SI Fig. S3, SI Materials and Methods, section B).

The best FRSM model included MAT and MAP as climate variables to the detriment of other combinations (SI Table S7). They provided non-correlated information about the climatic characteristics of the training trials ($r=-0.11$; $p=0.78$). Based on this model, response surfaces to climate were obtained at the ecotype level, and for the whole dataset (i.e., general performance model; see SI Materials and Method, section B).

Future forecasts of climate responses

For the current distribution area of *P. halepensis* (EUFORGEN, 2009), height growth models were projected at 30s arcsecond resolution (*ca.* 1 km²) using climatic maps for ‘reference’ conditions (i.e., the period 1961-1990 AD) from WorldClim (Hijmans *et al.*, 2005) and for ‘future’ conditions (i.e., the period 2071-2100 AD). For future conditions, we used the climate projections from Sáenz-Romero *et al.* (2017) under two representative concentration pathways: a ‘moderate’ scenario (RCP 4.5 watts per m⁻²; hereafter RCP 4.5) and a ‘pessimistic’ scenario (RCP 8.5 watts per m⁻²; hereafter RCP 8.5). Based on the general performance model (i.e., disregarding ecotypic variation), and also for each of the ecotypic models independently, we projected tree height for every pixel of the current distribution of *P. halepensis*. We then produced maps of height differences between reference conditions and height projections, as a percentage of reference height of each pixel (*ca.* 306,000 occurrence points; EUFORGEN, 2009). Maps were generated under the RCP 4.5 and 8.5 scenarios, and for the general performance and each ecotype model (see SI Materials and Methods, section C). To delineate the spatial patterns of ecotypic performance, we compared the relevance of differential plasticity on tree height. We first assumed that each ecotype could be potentially distributed over the entire distribution range of the species (hereafter, regional distribution). In this way, we made the comparison easier among ecotypes, and therefore, the implication of these results better assessed for management purposes (e.g., deployment or assisted migration strategies).

We then produced a map of the ‘tallest ecotype’ by projecting the ecotype with the highest height for every pixel of the distribution and under each scenario. This map was filtered at the pixel level for tree height of the tallest ecotype, which exceeded the second-best ecotype by at least 0.26 m (Least Significant Difference of ecotype means at $p=0.05$ in model [1]). Moreover, we created histograms to summarize the height frequency distribution under reference and future climates for general performance and each ecotype separately and independently for (i) regional distribution (i.e., maximum occupancy) and (ii) for local conditions (i.e., local occupancy). Non-parametric statistics were used to assess changes in frequency distribution (local *vs.* regional, reference *vs.* future climate; SI Materials and Methods, section C).

Results

Evidence of differential ecotypic plasticity linked to climate

The thermophilic *P. halepensis* showed phenotypic plasticity for the total height that was genetically structured, as revealed by significant ecotype by environment interactions (Likelihood-Ratio test = 15, $p < 0.001$, for models with and without ecotype by trial interaction term). According to the selected factorial regression response surface model (FRSM), the nature and magnitude of differential tree height responses were adequately described by simple climate factors, mean annual temperature (MAT) and mean annual precipitation (MAP) in particular (marginal $R^2 = 43.1\%$; SI Table S7). For reference climate conditions (period 1961-1990 AD), and ignoring ecotypic variation, tree height at age 15 years exhibited a 2.5-fold increase with increased precipitation and temperature (from 4 to 10 m approximately) and with the largest tree height mainly found for locations with $MAP > 75\%$ of potential evapotranspiration (Fig. 1a). FRSM, however, pointed to variable ecotypic responses to precipitation (MAP, $MAP \times MAP$) and its combined effect with temperature ($MAT \times MAP$) (SI Table S8), and allowed quantification of specific sensitivities to climate for each ecotype (Table S9). These sensitivities produced height variability underlying $G \times E$ interactions (i.e., differential ecotypic plasticity) and also ecotypic differentiation in mean height across trials (i.e., standing ecotypic variation, with up to 13% relative difference between extreme ecotypes; Table 1). The FRSM-derived ecotypic equations (Table 1), which are solved in Fig. 1 for the climate envelope defined by the training trials, had, in most cases, coefficients of determination equal or higher to 0.75, indicating a reasonable fit. The relationship between tree height and climate, therefore, varied if ecotypes were considered (Fig. 1b-f). The standard deviations of predicted tree height, obtained through bootstrap of regression estimates, are shown in Fig. 2. Most ecotype predictions had deviations below 0.3 m across MAT and MAP combinations, but the weakest prediction power corresponded to ecotype DHT, followed by WHC. Also, the weakest spots for prediction purposes across ecotypes corresponded to the four combinations of extreme MAT and MAP values, especially those of high MAP (Fig. 2).

The MAP variation across trials (from 300 to 800 mm) underlay varying ecotypic height performances. The Wet-summer/sub-Humid/Cool (WHC) ecotype (mainly from NE Spain and SE France) displayed the sharpest dependence on precipitation, as indicated by significant above- (linear) and below-average (quadratic) sensitivities to MAP (Table S9). Conversely, the Dry-summer/Semiarid/Temperate (DST) ecotype (from coastal SE Spain, inland Algeria and Tunisia, and Israel) and the Dry-summer/sub-Humid/Temperate (DHT) ecotype (from Greece and coastal Tunisia)

showed a subtle sensitivity to precipitation, for either linear (DST, DHT) or quadratic (DHT) sensitivities (Table S9). Contrary to precipitation, responses to temperature were approximately uniform across ecotypes (SI Table S8) and denoted a positive reaction to warming for the whole species (Table 1). In some instances, however, the abovementioned responses to MAP were contingent on the simultaneous increase in MAT and MAP (Table S9), which strongly benefitted DHT while penalizing WHC under the same conditions. Based on the ecotypic equations, the combined sensitivities to climate factors translated into height differences between extreme ecotypes (DHT and DSC) of 0.8 m at the shortest trial (Ademuz, Spain) and 2 m at the tallest trial (Oria, Italy) (SI Table S1).

Cross-validation on the training dataset (Table 1) indicated that the performance of ecotypic models was reasonably good, with mean absolute errors of *ca.* 1 m (height range of population-trial combinations = 10.2 m) and with correlations equal or higher to 0.75 between observed and predicted values. Using independent validation, the predictive ability of ecotypic models was superior (WST) or similar (DSC, DST, WHC) to that of the general performance model (ASE statistic; Table 1). The only exception was DHT, whose heights were most often underestimated by both the ecotypic and general models (SI Table S10). Overall, these results demonstrated the good predictive ability of climate functions.

Description of ecotypic responses to climate

The optimum height trajectories for the current average climate of each ecotype (curved black arrows in Fig. 1) indicated the direction and strength of maximum response and provided information on differential plasticity. In particular, DST responded positively to MAT up to a mean temperature of *ca.* 17°C, beyond which it showed a mild sensitivity to MAP (Fig. 1b). In contrast, the Dry-summer/Semiarid/Cold (DSC) ecotype (from inland Spain and Morocco) showed positive responses to increasing temperature (up to 1-2°C beyond their current mean climate) when accompanied by increasing precipitation (Fig. 1c). DHT showed positive, strong responses to temperature and precipitation (Fig. 1d), attaining the maximum height of all ecotypes within its climate envelope (≈ 12 m). Finally, the Wet-summer/Semiarid/Temperate (WST) ecotype (from the Balearic Islands, Italy, and E Spain) and the WHC ecotype – with the coldest and wettest origin – mildly benefitted from increased temperature and

precipitation (Fig. 1e,f). Most ecotypes displayed saddle-shaped response surfaces (Fig. 1), with no obvious combination of temperature and precipitation being optimum for tree height (SI Table S9). The exception was WHC, projected to reach a climate optimum at 1500 mm MAP, which is well above its current local climate at SE France and NE Spain (SI Table S9).

Spatial patterns of current performance

The climate scope of populations at their origin matched well the species' current climate envelope, as shown in Fig. 3a, left panel (average climate differences between midpoint of populations and midpoint of species distribution range: MAT = +1.3 °C, MAP = -62 mm). Comparing the differential plasticity of ecotypes to climate, DHT and WHC should attain, at present, the highest and lowest median height respectively for the entire region (M_R = 6.82 m and 6.58 m for DHT and WHC; Fig. 4). These results resembled the ecotypic ranking across trials (Table 1), which supported the effectiveness of the training trials for modeling purposes. We could also observe that DHT and WHC should perform best, at present, in extensive areas of the Mediterranean basin (Fig. 3a, central panel). We observed much larger median differences among ecotypes locally, i.e., considering only locations of each ecotype at its present occurrence. These differences amounted to *ca.* 3 m (M_L values, Fig. 4). Indication of local adaptation was limited to the mesic ecotypes DHT and WHC, which often showed a significantly higher height at origin (>40% of their local area) when compared with the rest of ecotypes (Fig. 3a, right panel).

Spatial forecasts under future conditions

Projections for the period 2071-2100 AD indicated a widespread increase in tree height across the species' current distribution (Fig. 5; more details in SI Figs. S4-S7 & Table S11). This increase could occur although climate change is projected to push most ecotypes beyond their current temperature range for both RCP 4.5 (1.8-3.5°C warmer and 5-111 mm drier on average) and 8.5 (3.6-5.7°C warmer and 73-188 mm drier on average) (Figs. 3b,c, left panels, and SI Table S12). However, wet summer ecotypes were the exception, for which growth inhibition was often projected both regionally (WHC) and locally (WST and WHC).

For RCP 4.5, the general performance model, disregarding ecotypic variation, projected height enhancement in 51% of the current species' distribution (N, NW, and

SE Mediterranean, and continental areas of northern Africa; Fig. 5a). However, notable differences were observed among the ecotypes (Fig. 5). Regionally, the largest area of height enhancement (65%) was projected for DHT, followed by DST (59%). Inhibition at over 40% of the distribution was projected for WHC (see also SI Fig. S6a). Overall, height inhibition of up to 1.4 m and enhancement of up to 2 m was expected (5% and 95% quantiles of height differences, respectively, for general performance; SI Fig. S6a, Fig. S7a). Locally, dry summer ecotypes (DST, DSC, and DHT) were projected to respond positively to warming, whereas wet summer ecotypes (WST and WHC mainly to experience stability or inhibition (Fig. 5). These responses translated into significant height differences among ecotypes both regionally (difference in median values between extreme ecotypes = 1.4 m) and locally (SI Fig. S4). Locally, however, differences were reduced from *ca.* 3 m (for reference conditions) to 1.7 m, and the ecotypes showing superiority at origin changed compared with reference conditions, with the growth of DHT being favored and WHC inhibited under moderate warming (Fig. 3b, right panel).

For RCP 8.5, the general performance model projected height enhancement for a slightly smaller fraction of the current distribution (48%), compared to RCP 4.5 (Fig. 5b). Overall, height inhibition of up to 2.8 m and enhancement of up to 2.5 m was expected (5% and 95% quantiles of height differences, respectively, for general performance; SI Fig. S6b, Fig. S7b). The dry summer DST and DSC ecotypes still showed superior growth compared to under reference conditions (Fig. 5), both regionally and locally. Conversely, DHT and WST ecotypes mainly experienced enhancement regionally, and WHC largely exhibited both regional and local inhibition (Fig. 5b). Only the dry summer DSC and DHT ecotypes exhibited superiority at origin (Fig. 3c, right panel). These contrasting responses exacerbated the height differences among ecotypes projected for RCP 4.5 regionally (difference in median values between extreme ecotypes = 2.3 m), but not locally (difference = 0.7 m; SI Fig. S5). While DST, DSC, and WST showed a significantly higher median height regionally than locally under both climate scenarios, the reverse situation was observed for ecotypes DHT and WHC (SI Fig. S4, Fig. S5).

From these results, it could be deduced that climate change would lead to smaller height differences among ecotypes, assuming there is local occupancy (i.e., no dispersal), especially for the most stringent RCP 8.5 scenario. This was mainly because the dry summer ecotypes benefitted from warming at their origin. Conversely, height

differences were amplified under maximum occupancy. Accordingly, the climate models forecasted that WHC would be outperformed by DHT and also by DSC, across the whole distribution range (Fig. 3b,c).

Discussion

We developed, for the widespread thermophilic conifer *Pinus halepensis*, ecotype-specific vertical growth responses to climate which were then projected to reference and future climate conditions (2071-2100 AD) across the Mediterranean. Tree height inferences based on a factorial regression response surface model (FRSM) revealed good predictive ability of ecotypic climate functions. Importantly, the array of provenance trials used closely approximated the species' climate envelope, which is critical for establishing realistic projections (Wang *et al.*, 2006). Additionally, the climate-based ecotypic models were validated using an independent dataset, which reduced the uncertainty in our projections of climate responses for the species. Our approach, based on ecotypic classification, can provide parsimonious and, thus, hands-on information on the adaptive structure of a species to forest managers and end-users of genetic material. Other methods to investigate intraspecific variability may cluster populations based directly on their trait responses (DeLacy *et al.*, 1996), or fit 'universal response functions' as an interaction between the trial's climate and the climate of a population's origin (Wang *et al.*, 2010). However, these methods are sensitive to unbalanced data and may require intricate procedures for estimating population performance at sites with extreme climate (Wang *et al.*, 2010).

Despite the importance of gene flow in *P. halepensis* (Steinitz *et al.*, 2012), the FRSM model indicated that differentiation by climate is strong throughout the entire species' range (Aitken & Bemmels, 2015). Simple climate factors such as mean annual temperature and precipitation adequately described the nature and magnitude of intraspecific vertical growth responses. Therefore, when ecotypes are considered, the relationship between tree height and climate varies considerably. However, evidence of local adaptation was limited to mesic ecotypes. For the remaining ecotypes, the absence of local adaptation for tree height could be related to complex trade-offs among traits maximizing fitness for this species (Santini *et al.*, 2019), or to the existence of adaptation lags related to historical climate (e.g., Rehfeldt *et al.*, 2002). Importantly, the climate-based grouping of ecotypes matched faithfully a recent definition of genetic groups of the species based on molecular (putatively neutral) SNP information (Serra-

Varela *et al.*, 2017), which identified populations with similar evolutionary histories. The observed match between classifications is supportive (but not conclusive) of the impact of selection processes under different environmental conditions, as some SNP markers used by Serra-Varela *et al.* (2017) were likely influenced by adaptive selection. As such, our results suggest that selection by climate may have favored different genotypes in populations subjected to distinctive bottlenecks during the post-glacial westward Mediterranean expansion of *P. halepensis* (Grivet *et al.*, 2009).

Indeed, the relevance of climate in shaping adaptive patterns for *P. halepensis* has been well recognized for many morpho-physiological traits, including growth and reproduction (e.g. Calamassi *et al.*, 2001; Climent *et al.*, 2008; Schiller & Atzmon, 2009; Klein *et al.*, 2013; Voltas *et al.*, 2015; David-Schwartz *et al.*, 2016). Altogether, our results support range-wide GxE regarding climate in *P. halepensis* (Savolainen *et al.*, 2007; Peterson *et al.*, 2019). This thermophilic pine is known to exhibit a general adaptive syndrome by which less plastic populations are characterized by higher survival and larger investment in reproduction and reserves than in growth (Climent *et al.*, 2008; Voltas *et al.*, 2018; Santini *et al.*, 2019). In particular, intraspecific variation in tree height was mainly associated with precipitation, suggesting that variation in natural selection is more related to water availability than to temperature. However, this is not unexpected for a conifer subject to steep gradients of water availability (Richardson *et al.*, 2014). Therefore, the observed height responses, deduced from ecotype-specific climate sensitivities, are fundamental for interpreting intraspecific adaptation (Benito-Garzón & Fernández-Manjarrés, 2015).

The projected climate responses presented an evident northeast-southwest cline with mesic ecotypes outperforming ecotypes of drier climates across a large proportion of the species' distribution area. In particular, it indicated that ecotypes WHC and DHT should grow best, at present, in extensive areas of the Mediterranean basin. These ecotypes from northern and eastern Mediterranean areas allocate fewer resources to reproduction (Climent *et al.*, 2008) and transpire more in summer (Santini *et al.*, 2019), but they do not show reduced survival than their drier counterparts from southern and western regions (Voltas *et al.*, 2018). Particularly, DHT (native to the eastern Mediterranean basin) bears high genetic diversity and, importantly, encompasses most of the functional variation of *P. halepensis* (Grivet *et al.*, 2009). Along with this suite of features, the large climate-related height plasticity of DHT suggests that this ecotype may be decisive for the persistence of species through climate change and, thus,

important for the implementation of deployment or assisted migration strategies. Although taller trees may be more vulnerable to conduction-blocking embolisms (Olson *et al.*, 2018) and, therefore, to warming-induced drought stress, David-Schwartz *et al.* (2016) have shown that eastern populations of the species are not more embolism-susceptible than their western counterparts (David-Schwartz *et al.*, 2016). The latter suggests that fast vertical growth and drought tolerance are compatible within *P. halepensis*.

The larger height differences observed locally than range-wide among ecotypes could be mainly a consequence of more favorable conditions occurring at the wettest spots of the species' range for a conifer limited by drought stress (Sáenz-Romero *et al.*, 2019). However, those height differences at the local level also reflected differential ecotypic sensitivities to climate, as revealed in this study. Notably, the lack of superiority at the origin for most ecotypes suggests that differential plasticity is not necessarily locally adaptive, at least for vertical growth, despite the existence of intraspecific plastic variation for tree species with long lifespans (Benito-Garzón *et al.*, 2019; Gárate-Escamilla *et al.*, 2019). In this regard, while specialization in harsh environments is likely linked to limited plasticity and a conservative resource-use strategy, the opposite is to be expected for specialization in favorable environments (Valladares *et al.*, 2007).

Under future climate (i.e., 2071-2100 AD) and despite the projected warmer and drier conditions, projections indicated widespread height enhancement of *P. halepensis*, but smaller height differences among ecotypes assuming there is local occupancy (i.e., no dispersal), especially for the most stringent RCP 8.5 scenario. Height enhancement becomes particularly evident for the most xeric populations of the western Mediterranean, which are forecasted to be favored even under extreme warming. On the other hand, the projected height inhibition of wet summer ecotypes, either regionally (WHC) or locally (WST and WHC), suggests that these ecotypes could be at risk. The future height decline of wet summer ecotypes may lead to forest fragmentation as a result of maladaptation to rising drought (Montwé *et al.*, 2016). Importantly, forecasted trends for RCP 4.5 and RCP 8.5 were similar, but with more inhibition under the more severe RCP 8.5 for wet summer ecotypes. Nevertheless, our dataset lacked test trials at sites warmer than 20°C MAT, which represents 17% and 37% of the distribution of the species under RCP 4.5 and 8.5, respectively. Thus, the reliability of projections beyond this temperature threshold might be compromised.

Prevalent positive responses to warming are supported by studies reporting radial growth stimulation of old Aleppo pine forests even in semi-arid/arid and warm environments such as the pre-Saharan Algerian steppe (Choury *et al.*, 2017) or Israel (Dorman *et al.*, 2015). These responses could involve a shift of the actual growing season towards earlier months in winter-spring (Bigelow *et al.*, 2014). This shift would make growth more reliant on water availability at the very early growing season (Choury *et al.*, 2017), a period progressively more suitable for tree performance as the climate becomes warmer and drier. Also, elevated CO₂ stimulation effects on growth cannot be discarded, although uniformity of ecotypic height responses to elevated CO₂ can be expected (Resco de Dios *et al.*, 2016). Anyhow, tree height responses to eCO₂ are indirectly accounted for in our projections (see SI Note on eCO₂ effects). Other studies, however, have reported drought-induced declines in radial growth for *P. halepensis* under severe dry spells (de Luis *et al.*, 2013; Gazol *et al.*, 2017). In this regard, weather extremes can have a substantial impact on Aleppo pine performance (Santos del Blanco *et al.*, 2013), which could only be approximated by the climate averages used in our study. The sequence of consecutive extreme weather events and, also, the dynamics of biotic damage will, almost certainly, tip the balance towards survival or mortality (Sangüesa-Barreda *et al.*, 2015), as future conditions will likely produce a mismatch between the species' fitness limits and its current distribution. On the other hand, our projections agree with alternative method-testing scenarios for assisted migration in *P. halepensis*, in which increased volume is projected for 2050 AD, except in SE France and coastal Spain (Benito-Garzón & Fernández-Manjarrés, 2015).

In our approach, we substituted space for time in forecasting height growth responses to climate (but see Merilä & Hendry, 2014). Nevertheless, the evolutionary history of *P. halepensis* also involves other fitness traits than height, and also trade-offs mainly related to drought and fire events (Aitken & Whitlock, 2013; Pausas, 2015; Santini *et al.*, 2019), that could be further used to describe adaptive patterns. Alternative methodologies that remain to be tested could involve tree-ring records and FRSMs by untangling the effects of the site (space) and year (time) in the explanation of differential ring-width responses in trial networks. Indeed, dendrochronological approaches applied to common-garden experiments are emerging that aim to forecast responses to climate fluctuations (Montwé *et al.*, 2015; George *et al.*, 2019; Suvanto *et al.*, 2016). Also, the combination of height records with radial growth modeling could

improve estimates of productivity and carbon sequestration potential (Klesse *et al.*, 2018). For a holistic perspective of our short-term projections, future studies should compare these forecasts to long-term dispersal rates and gene flows, which are considered to be high in *P. halepensis* (Steinitz *et al.*, 2012; Serra-Varela *et al.*, 2017).

This study demonstrates that phenotypic changes in tree height, for a widespread and ecologically diverse conifer such as *P. halepensis*, are, to a large extent, genetically based and not exclusively the result of uniform plastic responses. We show that this variation is related to climate and precipitation in particular, and indeed, accounting for intraspecific sensitivity to climate is a crucial step for forecasting tree responses (Peterson *et al.*, 2019). The latter is exemplified here for *P. halepensis*, a species native to warm, dry environments of substantial ecological, cultural and economic importance which may have to confront very different conditions in the near future. Changes in climate may, therefore, drive nonintuitive shifts in range, particularly affecting the edge of the species' range in wetter regions. We do not yet know to what extent plasticity of the integrated phenotype (or the exhibition of particular arrays of traits in a changing environment) may be crucial for the resilience and persistence of forests in the near future. Nevertheless, the patterns we report here are crucial towards a comprehensive understanding of prevalent climate-driven trade-offs. Further insights are urgently needed to conclusively predict the evolutionary response of this exemplary Mediterranean conifer to climate change and the reactions of forest trees typical of dry environments in general.

Acknowledgements

This study was funded by the Spanish Government (MINECO/FEDER coordinated grants AGL2015-68274-C3-3-R and RTI2018-094691-B-C3). We acknowledge the collaboration of the Spanish network of forest genetic trials (GENFORÉD) for technical cooperation, and E. Fusaro, F. Righi (CREA, Rome, Italy), G. Scarascia-Mugnozza and F. Riguzzi (University of Viterbo, Italy) for establishing the Italian trials and gathering data. We also thank G. Schiller (ARO, Volcani Center) for the establishment and maintenance of the provenance trials in Israel, together with the Forest Dept. of the KKL-JNF. The Italian trials were established under the project EU FAIR "Selection of Mediterranean Pines and Cedars for Sustainable Afforestation – MEDPINE") and the program FAO/SCM/CRFM/4bis.

639 **Author contribution:** JV, TK and GdM conceived and designed the study. TSP, TAS
640 and JV carried out the analyses. TSP and JV wrote the final draft of the manuscript. All
641 other authors contributed data and commented on the manuscript

For Peer Review

References

- Aitken SN, Whitlock MC. 2013. Assisted gene flow to facilitate local adaptation to climate change. *Annual Review of Ecology, Evolution, and Systematics* **44**: 367–88.
- Aitken SN, Bemmels JB. 2015. Time to get moving: assisted gene flow of forest trees. *Evolutionary Applications* **9**: 271–290.
- Alberto FJ, Aitken SN, Alía R, González-Martínez SC, Hänninen H, Kremer A, Lefèvre F, Lenormand T, Yeaman S, Whetten R, Savolainen O. 2013. Potential for evolutionary responses to climate change – evidence from tree populations. *Global Change Biology* **19**: 1645–1661.
- Benito-Garzón M, Fernández-Manjarrés JF. 2015. Testing scenarios for assisted migration of forest trees in Europe. *New Forests* **46**: 979–994.
- Benito Garzón M, Robson TM, Hampe A. 2019. Δ TraitSDMs: species distribution models that account for local adaptation and phenotypic plasticity. *New Phytologist* **222**, 1757–1765.
- Bigelow SW, Papaik MJ, Caum C, North MP. 2014. Faster growth in warmer winters for large trees in a Mediterranean-climate ecosystem. *Climatic Change* **123**: 215–224.
- Calamassi R, Della Roca G, Falusi M, Paoletti E, Strati S. 2001. Resistance to water stress in seedlings of eight European provenances of *Pinus halepensis* Mill. *Annals of Forest Science* **58**: 663–672.
- Choury Z, Shestakova TA, Himrane H, Touchan R, Kherchouche D, Camarero JJ, Voltas J. 2017. Quarantining the Sahara desert: growth and water-use efficiency of Aleppo pine in the Algerian Green Barrier. *European Journal of Forest Research* **136**: 139–152.
- Climent J, Prada MA, Calama R., Chambel MR, de Ron DS, Alía R. 2008. To grow or to seed: ecotypic variation in reproductive allocation and cone production by young

female Aleppo pine (*Pinus halepensis*, Pinaceae). *American Journal of Botany* **95**: 833–842.

David-Schwartz R, Paudel I, Mizrachi M, Delzon S, Cochard H, Lukyanov V, Bade E, Capdeville G, Shklar G, Cohen S. 2016. Indirect evidence for genetic differentiation in vulnerability to embolism in *Pinus halepensis*. *Frontiers in Plant Science* **7**: 768.

de Luis M, Čufar K, Di Filippo A, Novak K., Papadopoulos A, Piovesan G, Rathgeber CBK, Raventós J, Saz MA, Smith KT. 2013. Plasticity in dendroclimatic response across the distribution range of Aleppo pine (*Pinus halepensis*). *PLoS One* **8**: e83550.

DeLacy I, Basford KE, Cooper M, Bull JK, McLaren CG. 1996. Analysis of multi-environment trials – An historical perspective. In: M. Cooper & G.L. Hammer (Eds.), *Plant Adaptation and Crop Improvement* (pp. 39-124), Wallingford, UK: CAB International.

Denis JB, Piepho HP, van Eeuwijk FA. 1997. Modelling expectation and variance for genotype by environment data. *Heredity* **79**: 162–171.

Dorman M, Svoray T, Perevolotsky A, Moshe Y, Sarris D. 2015. What determines tree mortality in dry environments? A multi-perspective approach. *Ecological Applications* **25**: 1054-1071.

Esteban LG, Martín JA, de Palacios P, García-Fernández F, López R. 2010. Adaptive anatomy of *Pinus halepensis* trees from different Mediterranean environments in Spain. *Trees* **24**: 19–30.

EUFORGEN. 2009. Distribution map of Aleppo pine (*Pinus halepensis*). Available at: www.euforgen.org [Accessed May 6, 2019].

- 710 **Fréjaville T, Vizcaíno-Palomar N, Fady B, Kremer A, Benito Garzón M.** 2020.
 711 Range margin populations show high climate adaptation lags in European trees. *Global*
 712 *Change Biology* **26**: 484–495.
 713
- 714 **Gárate-Escamilla H, Hampe A, Vizcaíno-Palomar N, Robson TM, Benito-Garzón**
 715 **M.** 2019. Range-wide variation in local adaptation and phenotypic plasticity of
 716 fitness-related traits in *Fagus sylvatica* and their implications under climate change.
 717 *Global Ecology and Biogeography* **28**: 1336–1350.
 718
- 719 **Gazol A, Ribas M, Gutiérrez E, Camarero JJ.** 2017. Aleppo pine forests from across
 720 Spain show drought-induced growth decline and partial recovery. *Agricultural and*
 721 *Forest Meteorology* **232**: 186–194.
 722
- 723 **George JP, Grabner M, Campelo F, Karanitsch-Ackerl S, Mayer K., Klumpp RT,**
 724 **Schüler S.** 2019. Intra-specific variation in growth and wood density traits under water-
 725 limited conditions: Long-term-, short-term-, and sudden responses of four conifer tree
 726 species. *Science of the Total Environment* **660**: 631–643.
 727
- 728 **Grivet D, Sebastiani F, González-Martínez SC, Vendramin GG.** 2009. Patterns of
 729 polymorphism resulting from long-range colonization in the Mediterranean conifer
 730 Aleppo pine. *New Phytologist* **184**: 1016–1028.
 731
- 732 **Harris I, Jones PD, Osborn TJ, Lister DH.** 2014. Updated high-resolution grids of
 733 monthly climatic observations – the CRU TS3.10 Dataset. *International Journal of*
 734 *Climatology* **34**: 623–642.
 735
- 736 **Heslot N, Akdemir D, Sorrells ME., Jannink JL.** 2014. Integrating environmental
 737 covariates and crop modeling into the genomic selection framework to predict genotype
 738 by environment interactions. *Theoretical and Applied Genetics* **127**: 463–480.
 739
- 740 **Hijmans RJ, Cameron SE, Parra JL, Jones PG, Jarvis A.** 2005. Very high resolution
 741 interpolated climate surfaces for global land areas. *International Journal of Climatology*
 742 **25**: 1965–1978.
 743

- Kawecki TJ, Ebert D.** 2004. Conceptual issues in local adaptation. *Ecology Letters* **7**: 1225–1241.
- Klein T, Di Matteo G, Rotenberg E, Cohen S, Yakir D.** 2013. Differential ecophysiological response of a major Mediterranean pine species across a climatic gradient. *Tree Physiology* **33**: 26–36.
- Klesse S, Babst F, Lienert S, Spahni R, Joos F, Bouriaud O, Carrer M, Di Filippo A, Poulter B, Trotsiuk V, Wilson R, Frank DC.** 2018. A combined tree ring and vegetation model assessment of European forest growth sensitivity to interannual climate variability. *Global Biogeochemical Cycles* **32**: 1226–1240.
- Le Houérou HN.** 2004. An agro-bioclimatic classification of arid and semiarid lands in the isoclimatic Mediterranean zones. *Arid Land Research and Management* **18**: 301–346.
- Malosetti M., Ribaud JM, van Eeuwijk FA.** 2013. The statistical analysis of multi-environment data: modeling genotype-by-environment interaction and its genetic basis. *Frontiers in Physiology* **4**: 44.
- Mátyás C.** 1996. Climatic adaptation of trees: rediscovering provenance tests. *Euphytica* **92**: 45–54.
- Mátyás C, Nagy L, Jármai EU.** 2009. Genetic background of responses of trees to aridification at the xeric forest limit and consequences for bioclimatic modelling. In: K. Strělcová *et al.* (Eds.), *Bioclimatology and Natural Hazards* (pp. 179-196), Dordrecht: Springer Science & Business Media.
- Merilä J, Hendry AP.** 2014. Climate change, adaptation, and phenotypic plasticity: the problem and the evidence. *Evolutionary Applications* **7**: 1–14.
- Moles AT, Warton DI, Warman L, Swenson NG, Laffan SW, Zanne AE, Pitman**

- 778 **A, Hemmings FA, Leishman MR.** 2009. Global patterns in plant height. *Journal of*
779 *Ecology* **97**: 923-932.
- 780
- 781 **Montwé D, Spiecker H, Hamann A.** 2015. A five decades of growth in a genetic field
782 trial of Douglas-fir reveal trade-offs between productivity and drought tolerance. *Tree*
783 *Genetic and Genomes* **11**: 29.
- 784
- 785 **Montwé D, Isaac-Renton M, Hamann A, Spiecker H.** 2016. Drought tolerance and
786 growth in populations of a wide-ranging tree species indicate climate change risks for
787 the boreal north. *Global Change Biology* **22**: 806–815.
- 788
- 789 **Nakagawa S, Schielzeth H.** 2013. A general and simple method for obtaining R^2 from
790 generalized linear mixed-effects models. *Methods in Ecology and Evolution* **4**: 133–
791 142.
- 792
- 793 **Neale DB, Wheeler NC.** 2019. Adaptive genetic variation. In: D. B. Neale & N. C.
794 Wheeler (Eds.), *The Conifers: Genomes, Variation and Evolution* (pp. 225–254). New
795 York, NY: Springer Science & Business Media.
- 796
- 797 **Ne’eman G, Goubitz S, Nathan R.** 2004. Reproductive traits of *Pinus halepensis* in
798 the light of fire – a critical review. *Plant Ecology* **171**: 69–79.
- 799
- 800 **Olson ME, Soriano D, Rosell JA, Anfodillo T, Donoghue MJ, Edwards EJ, León-**
801 **Gómez C, Dawson T, Camarero JJ, Castorena M, Echeverría A, Espinosa CI,**
802 **Fajardo A, Gazol A, Isnard S, Lima RS, Marcati CR, Méndez-Alonzo R.** 2018.
803 Plant height and hydraulic vulnerability to drought and cold. *Proceedings of the*
804 *National Academy of Sciences* **115**: 7551-7556.
- 805
- 806 **O’Neill GA, Hamann A, Wang T.** 2008. Accounting for population variation improves
807 estimates of the impact of climate change on species’ growth and distribution. *Journal*
808 *of Applied Ecology* **45**: 1040–1049.
- 809
- 810 **Pausas JG.** 2015. Evolutionary fire ecology: lessons learned from pines. Trends in
811 *Plant Science* **20**: 318–324.

- Peterson M, Doak DF, Morris WF.** 2019. Incorporating local adaptation into forecasts of species' distribution and abundance under climate change. *Global Change Biology* **25**: 775–793.
- Rehfeldt GE, Tchebakova NM, Parfenova YI, Wykoff WR., Kuzmina NA, Milyutin LI.** 2002. Intraspecific responses to climate in *Pinus sylvestris*. *Global Change Biology* **8**: 912–929.
- Reich PB, Oleksyn J.** 2008. Climate warming will reduce growth and survival of Scots pine except in the far north. *Ecology Letters* **11**: 588–597.
- Resco de Dios V, Mereed TE, Ferrio JP, Tissue DT, Voltas J.** 2016. Intraspecific variation in juvenile tree growth under elevated CO₂ alone and with O₃: a meta-analysis. *Tree Physiology* **6**: 682–693.
- Richardson JL, Urban MC, Bolnick DI, Skelly DK.** 2014. Microgeographic adaptation and the spatial scale of evolution. *Trends in Ecology & Evolution* **29**: 165–173.
- Sáenz-Romero C, Lamy JB, Ducousso A, Musch B, Ehrenmann F, Delzon S, Cavers S, Chalupka W, Dağdaş S, Hansen JK, Lee SJ, Liesebach M, Rau HM, Psomas A, Schneck V, Steiner W, Zimmermann NE, Kremer A.** 2017. Adaptive and plastic responses of *Quercus petraea* populations to climate across Europe. *Global Change Biology* **23**: 2831–2847.
- Sáenz-Romero C, Kremer A, Nagy L, Ujvári-Jármay E, Ducousso A, Kóczán-Horváth A, Hansen JK, Mátyás C.** 2019. Common garden comparisons confirm inherited differences in sensitivity to climate change between forest tree species. *PeerJ* **7**: e6213.
- Sangüesa-Barreda G, Linares JC, Camarero JJ.** (2015) Reduced growth sensitivity to climate in bark-beetle infested Aleppo pines: Connecting climatic and biotic drivers of forest dieback. *Forest Ecology and Management* **357**: 126–137.

- 846
- 847 **Santini F, Climent J, Voltas J.** 2019. Phenotypic integration and life history strategies
 848 among populations of *Pinus halepensis*: an insight through structural equation
 849 modeling. *Annals of Botany* **124**: 1161–1171.
- 850
- 851 **Santos del Blanco L, Bonser SP, Valladares F, Chambel MR, Climent J.** 2013.
 852 Plasticity in reproduction and growth among 52 range-wide populations of a
 853 Mediterranean conifer: adaptive responses to environmental stress. *Journal of*
 854 *Evolutionary Biology* **26**: 1912–1924.
- 855
- 856 **Savolainen O, Pyhäjärvi T, Knürr T.** 2007. Gene flow and local adaptation in trees.
 857 *Annual Review of Ecology, Evolution, and Systematics* **38**: 595–619.
- 858
- 859 **Sbay H, Zas R.** 2018. Geographic variation in growth, survival, and susceptibility to
 860 the processionary moth (*Thaumetopoea pityocampa* Dennis & Schiff.) of *Pinus*
 861 *halepensis* Mill. and *P. brutia* Ten.: results from common gardens in Morocco. *Annals*
 862 *of Forest Science* **75**: 69.
- 863
- 864 **Schiller G, Atzmon N.** 2009. Performance of Aleppo pine (*Pinus halepensis*)
 865 provenances grown at the edge of the Negev desert: a review. *Journal of Arid*
 866 *Environments* **73**: 1051–1057.
- 867
- 868 **Serra-Varela MJ, Alía R, Ruiz-Daniels R, Zimmermann NE, Gonzalo-Jiménez J,**
 869 **Grivet D.** 2017. Assessing vulnerability of two Mediterranean conifers to support
 870 genetic conservation management in the face of climate change. *Diversity &*
 871 *Distributions* **23**: 507–516.
- 872
- 873 **Sheil D.** 2018. Forests, atmospheric water and an uncertain future: the new biology of
 874 the global water cycle. *Forest Ecosystems* **5**: 19.
- 875
- 876 **Sixto H, Gil PM, Ciria P, Camps F, Cañellas L, Voltas J.** 2016. Interpreting
 877 genotype-by-environment interaction for biomass production in hybrid poplars under
 878 short-rotation coppice in Mediterranean environments. *Global Change Biology*
 879 *Bioenergy* **8**: 1124–1135.

- Skovsgaard JP, Vancley JK.** 2007. Forest site productivity: a review of the evolution of dendrometric concepts for even-aged stands. *Forestry* **81**: 13–31.
- Spencer DJ.** 1985. Dry country pines: provenance evaluation of the *Pinus halepensis* – *P. brutia* complex in the semi-arid region of south-east Australia. *Australian Forest Research* **15**: 263–279.
- Steinitz O, Robledo-Arnuncio JJ, Nathan R.** 2012. Effects of forest plantations on the genetic composition of conspecific native Aleppo pine populations. *Molecular Ecology* **21**: 300–313.
- Suvanto S, Nöjd P, Henttonen HM, Beuker E, Mäkinen H.** 2016. Geographical patterns in the radial growth response of Norway spruce provenances to climatic variation. *Agricultural and Forest Meteorology* **222**: 10–20.
- Tapias R, Climent J, Pardos JA, Gil L.** 2004. Life histories of Mediterranean pines. *Plant Ecology* **171**: 53–68.
- Valladares F, Gianoli E, Gómez JM.** 2007. Ecological limits to plant phenotypic plasticity. *New Phytologist* **176**: 749–763.
- Valladares F, Matesanz S, Guilhaumon F, Araújo MB, Balaguer L, Benito-Garzón M, Cornwell W, Gianoli E, van Kleunen M, Naya DE, Nicotra AB, Poorter H, Zavala MA.** 2014. The effects of phenotypic plasticity and local adaptation on forecasts of species range shifts under climate change. *Ecology Letters* **17**: 1351–1364.
- van Eeuwijk FA.** 1995. Linear and bilinear models for the analysis of multi-environment trials: I. An inventory of models. *Euphytica* **84**: 1–7.
- van Eeuwijk FA, Malosetti M, Yin X, Struik P, Stam P.** 2005. Statistical models for genotype by environment data: from conventional ANOVA models to eco-physiological QTL models. *Australian Journal of Agricultural Research* **56**: 883–894.

- Viglizzo EF, Jobbágy EG, Ricard MF, Paruelo JM.** 2016. Partition of some key regulating services in terrestrial ecosystems: Meta-analysis and review. *Science of The Total Environment* **562**: 47-60.
- Voltas J, Chambel MR, Prada MA, Ferrio JP.** 2008. Climate-related variability in carbon and oxygen stable isotopes among populations of Aleppo pine grown in common-garden tests. *Trees* **22**: 759–769.
- Voltas J, Lucabaugh D, Chambel MR, Ferrio JP.** 2015. Intraspecific variation in the use of water sources by the circum-Mediterranean conifer *Pinus halepensis*. *New Phytologist* **208**: 1031–1041.
- Voltas J, Shestakova TA, Patsiou T, di Matteo G, Klein T.** 2018. Ecotypic variation and stability in growth performance of the thermophilic conifer *Pinus halepensis* across the Mediterranean basin. *Forest Ecology and Management* **424**: 205–215.
- Wang T, Hamann A, Yanchuk A, O'Neill GA, Aitken SN.** 2006. Use of response functions in selecting lodgepole pine populations for future climates. *Global Change Biology* **12**: 2404–2416.
- Wang T, O'Neill G, Aitken SN.** 2010. Integrating environmental and genetic effects to predict responses of tree populations to climate. *Ecological Applications* **20**: 153-163.
- Ward JR.** 1963. Hierarchical grouping to optimize an objective function. *Journal of the American Statistical Association* **58**: 236–44.
- Williams GC.** 1966. Adaptation and natural selection. Princeton: Princeton University Press.
- Zavala MA, Espelta JM, Caspersen J, Retana J.** 2011. Interspecific differences in sapling performance with respect to light and aridity gradients in Mediterranean pine–oak forests: implications for species coexistence. *Canadian Journal of Forest Research* **41**: 1432–1444.

948 **Zhu K, Zhang J, Niu S, Chu C, Luo Y.** 2018. Limits to growth of forest biomass
949 carbon sink under climate change. *Nature Communications* **9**: 2709.
950

For Peer Review

Supporting Information content

Table S1. Geographic and climatic characteristics of the nine trials used as training dataset and of the five trials used as test dataset.

Table S2. Mean survival, number of populations per ecotype at each trial, characteristics of the experimental design, and layout of the nine trials used as training dataset and of the five trials used as test dataset.

Table S3. Geographical and climatic (CRU TS 3.24, 1961-1990 period; 4) characteristics of the 82 populations of *Pinus halepensis* used as the training dataset for factorial regression modeling.

Table S4. Main climate characteristics of ecotypes ($\pm$ standard deviation of population means) based on the WorldClim dataset.

Fig. S1. The distribution range of *Pinus halepensis* identified with hatching.

Fig. S2. Dendrogram of the 82 populations of *Pinus halepensis* used as the training dataset for factorial regression modeling of tree height at age 15.

SI Materials and Methods:

A. Study sites and populations assignment to ecotypes

B: Ridge analysis and validation of response surface functions

C. Quantification of ecotypic responses to future climate

Table S5. Information criteria used to select the best classification of populations into ecotypes for tree height at age 15.

Table S6. Probability of assignment to each ecotype of 44 new populations evaluated in test trials using linear discriminant functions.

Table S7. Factorial regression response surface models of tree height at age 15 partitioning the ecotype $\times$ trial interaction term.

Table S8. Full factorial regression response surface model (FRSM) of tree height at age 15 using mean annual temperature (MAT) and mean annual precipitation (MAP) at the trial level.

Table S9. Ecotypic sensitivities (β_{MAP} , $\beta_{\text{MAT} \times \text{MAP}}$, β'_{MAP}) to selected response surface terms of equation [5] of the main text related to mean annual temperature (MAT) and mean annual precipitation (MAP) for tree height (m) at age 15.

Table S10. Predictive performance of quadratic models at the ecotype level for tree height at age 15 of *Pinus halepensis* based on mean annual temperature and mean annual precipitation.

Table S11. Comparison of frequency distributions of tree height between reference and future climate conditions (RCP 4.5 and RCP 8.5) at the species (General Performance) and ecotype levels.

Table S12. Summary of climatic variables (MAT = mean annual temperature; MAP = mean annual precipitation) within the distribution of *Pinus halepensis* for reference and predicted future conditions under future RCP 4.5 and RCP 8.5 scenarios.

Fig. S3. Plot of predicted values versus residuals of the factorial regression response surface model, including mean annual temperature and mean annual precipitation.

Fig. S4. Spatial projections (future RCP 4.5 scenario) for tree height (m) at age 15 at the species (General Performance) (a) and ecotype level (b-f) for *Pinus halepensis* in the Mediterranean basin assuming maximum occupancy.

Fig. S5. Spatial projections (future RCP 8.5 scenario) for tree height (m) at age 15 at the species (General Performance) (a) and ecotype level (b-f) for *Pinus halepensis* in the Mediterranean basin assuming maximum occupancy.

Fig. S6. Cumulative distribution (frequency classes, left panels) and density plots (right panels) of the predicted height differences at age 15 between future and reference climate conditions.

Fig. S7. Tree height differences at age 15 between future and reference conditions in the Mediterranean basin.

Notes

Elevated CO₂ (eCO₂) levels effect on growth.

Fig. 1. Ecotypic responses of tree height (m) to mean annual temperature (MAT) and mean annual precipitation (MAP). (a) General performance (GP) of *Pinus halepensis* disregarding ecotypic variation, (b) Dry-summer/Semiarid/Temperate ecotype (DST), (c) Dry-summer/Semiarid/Cold ecotype (DSC), (d) Dry-summer/sub-Humid/Temperate ecotype (DHT), (e) Wet-summer/Semiarid/Temperate ecotype (WST), (f) Wet-summer/sub-Humid/Cool ecotype (WHC). Dots in (a) represent the climate conditions of the nine trials used for model building (training dataset). The X marks represent the

climate conditions of the five trials used for independent validation (test dataset). For each panel in (b-f), the cross represents the average climate conditions of the corresponding ecotype (and its 95% confidence ellipse, dashed area) based on WorldClim data (Hijmans *et al.*, 2005) for reference conditions (period 1961-1990 AD); the garnet red and bright red dots represent the forecasted climate (and standard deviation) for two representative concentration pathways, namely a ‘moderate’ scenario (RCP 4.5) and a ‘pessimistic’ scenario (RCP 8.5) respectively (note that the forecasted climate for ecotype DHT falls outside panel (f) and has estimates of 20.5 °C [MAT] and 541 mm [MAP]). The dashed lines in (a) depict isolines of fractional changes in the ratio of MAP to Potential Evapotranspiration (0.35 to 0.90), where PET is estimated as 70×MAT following Le Houérou (2004). The curved arrows represent the ridge of optimum response for changes in ± 15% height at age 15 years per ecotype (where a longer vector length indicates less plasticity).

Fig. 2. Standard deviation (STD) of predicted tree height responses to mean annual temperature (MAT) and mean annual precipitation (MAP) obtained from bootstrap of regression estimates via residual resampling. (a) General performance (GP) of *Pinus halepensis* disregarding ecotypic variation, (b) Dry-summer/Semiarid/Temperate ecotype (DST), (c) Dry-summer/Semiarid/Cold ecotype (DSC), (d) Dry-summer/sub-Humid/Temperate ecotype (DHT), (e) Wet-summer/Semiarid/Temperate ecotype (WST), (f) Wet-summer/sub-Humid/Cool ecotype (WHC). For each panel in (b-f), the cross represents the average climate conditions of the corresponding ecotype (and its 95% confidence ellipse, dashed area) based on WorldClim data (Hijmans *et al.*, 2005) for reference conditions (period 1961-1990 AD); the garnet red and bright red dots represent the forecasted climate (and standard deviation) for two representative concentration pathways, namely a ‘moderate’ scenario (RCP 4.5) and a ‘pessimistic’ scenario (RCP 8.5) respectively (note that the forecasted climate for ecotype DHT falls outside panel (f) and has estimates of 20.5 °C [MAT] and 541 mm [MAP]). The standard deviation ranges from 0.1 to 0.7 m, while most MAT-MAP combinations have a standard deviation < 0.3 m.

Fig. 3. Spatial projections of tallest ecotypes. Autoecology of *Pinus halepensis* distribution (*left*), spatial projections of tallest ecotypes for tree height (m) at age 15 years across the entire species’ distribution range in the Mediterranean basin,

considering maximum occupancy of each ecotype (*center*), and percentage of the area where each ecotype shows height superiority ($P < 0.05$) at the origin, i.e., under local occupancy (*right*). (a) Reference conditions (period 1961-1990 AD) based on WorldClim data (Hijmans *et al.*, 2005). (b) Future conditions (period 2071-2100 AD) using the climate layers for RCP 4.5. (c) Future conditions using the climate layers for RCP 8.5. In left panels, trials are depicted as black triangles (up triangles: training trials; down triangles: test trials) and populations as colored squares [populations and trials represented in (b) and (c) are based on reference climate]. In central panels, the pie charts represent the frequency of each ecotype significantly outperforming ($P < 0.05$) other ecotypes for each climate condition considering maximum occupancy. The species range is derived from the EUFORGEN distribution map (48) for a resolution of 10'. See the legend of Figure 1 for ecotype acronyms.

Fig. 4. Spatial projections for tree height (m) at age 15 under reference conditions (period 1961-1990 AD). (a) Species-level (General Performance) and (b, c, d, e, f) ecotype level, considering maximum occupancy in the Mediterranean basin. The histograms depict the class frequencies (%) of tree height at the ecotype and general level for the entire species' distribution. The class frequencies of tree height at the geographic origin of each ecotype, estimated for a 25 km radius around each ecotype's populations, are also shown in the histograms as hatched bars. A Median test (Z) for comparison between local (L) and regional (R) performance, together with their median values (M, in m), is included. Positive Z values denote local level is superior to regional level performance.

Fig. 5. Relative tree height differences at age 15 between future and reference conditions in the Mediterranean basin. *Left panels.* Maps of tree height differences (%) at the species level (General Performance, GP) for (a) RCP 4.5 scenario and (b) RCP 8.5 scenario between future (period 2071- 2100 AD) and reference conditions (period 1961-1990 AD) across the distribution range of *P. halepensis*. *Right panels.* Pie charts summarizing the relative importance of height differences (enhancement [$> 10\%$], inhibition [$< -10\%$], stability [$-10\%, +10\%$]) at the local (left charts) and regional (i.e., maximum occupancy of each ecotype, right charts) levels for general performance and for each ecotype. See the legend of Fig. 1 for ecotype acronyms and Fig. S7 in SI for absolute tree height differences.

1086 **Table 1.** Outline of ecotypic responses to climate factors. The number of population-trial combinations per ecotype (N), estimated ecotype means
1087 across trials (and their standard errors), full quadratic models (response surfaces) at the whole species (General Performance) and ecotype levels
1088 as a function of mean annual temperature (MAT) and mean annual precipitation (MAP), and predictive performance of ecotype models derived
1089 from a factorial regression response surface model for tree height at age 15 years. The predictive performance of ecotype models is evaluated by
1090 k -fold cross-validation based on $k = 10$ replicate samples (training dataset) and by the predictive ability of the response surfaces using
1091 independent data (test dataset). For cross-validation, the Mean Absolute Error (MAE) and the correlation (r) between observed and predicted
1092 values are shown. For predictive ability using independent data, the Average Squared Error (ASE , the sum of squared errors divided by the
1093 number of cases, N) of the corresponding ecotype model and the correlation (r) between observed and predicted values using the entire test
1094 dataset and a simplified dataset containing the subset of populations also evaluated in the training dataset (r^*) are shown. For comparison with
1095 ecotypic models, ASE values for the training and test datasets using the General Performance model are also presented. In the table, E_k is the
1096 fixed effect of the k^{th} ecotype, α (α') is the linear (quadratic) sensitivity of the whole species to a given environmental variable (or to the cross-
1097 product of variables MAT and MAP), and β (β') is the ecotype-specific linear (quadratic) sensitivity for the same variable (or cross-product of
1098 two variables MAT and MAP).
1099

1100

	Response surface equation										Cross-validation (train dataset)		Predictive ability (test dataset)			
	Mean height ¹ (m)	<i>N</i>	μ	α_{MAT}	α_{MAP}	$\alpha_{MAT \times MAP}$	α'_{MAT}	α'_{MAP}	R^2	<i>ASE</i>	<i>MAE</i> (m)	<i>r</i> (obs vs pred)	Ecotype model <i>ASE</i>	<i>r</i> (obs vs pred)	<i>r</i> [*] (obs vs pred)	GP model <i>ASE</i>
General Performance (GP)	-	248	-13.77	2.40	-0.00840	-0.00051	-0.05880	0.0000221	0.73	1.40	-	-	-	-	-	-
Ecotype			$\mu + E_k$	β_{MAT}	β_{MAP}	$\beta_{MAT \times MAP}$	β'_{MAT}	β'_{MAP}								
Dry-summer/Semiarid/ Temperate (DST)	6.71 (0.12)	64	-6.78	2.05	-0.02496	-0.00010	-0.05348	0.0000302	0.77	1.10	0.92	0.81	0.67	0.92	0.92	0.65
Dry-summer/Semiarid/ Cold (DSC)	6.53 (0.12)	72	-25.37	3.15	0.01623	-0.00168	-0.06502	0.0000151	0.78	0.90	0.71	0.87	0.52	0.96	0.98	0.61
Dry-summer/subHumid/ Temperate (DHT)	7.37 (0.15)	36	17.34	0.20	-0.06566	0.00132	-0.01703	0.0000502	0.67	2.55	1.30	0.83	3.67	0.86	0.90	2.89
Wet-summer/Semiarid/ Temperate (WST)	6.85 (0.13)	58	-21.92	2.85	0.00947	-0.00065	-0.07052	0.0000078	0.75	1.07	0.89	0.76	1.20	0.98	0.99	1.45
Wet-summer/subHumid/ Cool (WHC)	6.97 (0.23)	18	-28.44	3.51	0.01782	-0.00014	-0.10564	-0.0000051	0.75	1.21	1.24	0.75	0.26	0.97	-	0.13

¹ Ecotype least-square means obtained from equation [1]

1101

Ecotypic responses of tree height (m) to mean annual temperature (MAT) and mean annual precipitation (MAP). (a) General performance (GP) of *Pinus halepensis* disregarding ecotypic variation, (b) Dry-summer/Semi-arid/Temperate ecotype (DST), (c) Dry-summer/Semi-arid/Cold ecotype (DSC), (d) Dry-summer/sub-Humid/Temperate ecotype (DHT), (e) Wet-summer/Semi-arid/Temperate ecotype (WST), (f) Wet-summer/sub-Humid/Cool ecotype (WHC). Dots in (a) represent the climate conditions of the nine trials used for model building (training dataset). The X marks represent the climate conditions of the five trials used for independent validation (test dataset). For each panel in (b-f), the cross represents the average climate conditions of the corresponding ecotype (and its 95% confidence ellipse, dashed area) based on WorldClim data (Hijmans et al., 2005) for reference conditions (period 1961-1990 AD); the garnet red and bright red dots represent the forecasted climate (and standard deviation) for two representative concentration pathways, namely a 'moderate' scenario (RCP 4.5) and a 'pessimistic' scenario (RCP 8.5) respectively (note that the forecasted climate for ecotype DHT falls outside panel (f) and has estimates of 20.5 °C [MAT] and 541 mm [MAP]). The dashed lines in (a) depict isolines of fractional changes in the ratio of MAP to Potential Evapotranspiration (0.35 to 0.90), where PET is estimated as $70 \times \text{MAT}$ following Le

Houérou (2004). The curved arrows represent the ridge of optimum response for changes in $\pm 15\%$ height at age 15 years per ecotype (where a longer vector length indicates less plasticity).

186x239mm (300 x 300 DPI)

Standard deviation (STD) of predicted tree height responses to mean annual temperature (MAT) and mean annual precipitation (MAP) obtained from bootstrap of regression estimates via residual resampling. (a) General performance (GP) of *Pinus halepensis* disregarding ecotypic variation, (b) Dry-summer/Semi-arid/Temperate ecotype (DST), (c) Dry-summer/Semi-arid/Cold ecotype (DSC), (d) Dry-summer/sub-Humid/Temperate ecotype (DHT), (e) Wet-summer/Semi-arid/Temperate ecotype (WST), (f) Wet-summer/sub-Humid/Cool ecotype (WHC). For each panel in (b-f), the cross represents the average climate conditions of the corresponding ecotype (and its 95% confidence ellipse, dashed area) based on WorldClim data (Hijmans et al., 2005) for reference conditions (period 1961-1990 AD); the garnet red and bright red dots represent the forecasted climate (and standard deviation) for two representative concentration pathways, namely a 'moderate' scenario (RCP 4.5) and a 'pessimistic' scenario (RCP 8.5) respectively (note that the forecasted climate for ecotype DHT falls outside panel (f) and has estimates of 20.5 °C [MAT] and 541 mm [MAP]). The standard deviation ranges from 0.1 to 0.7 m, while most MAT-MAP combinations have a standard deviation < 0.3 m.

190x240mm (300 x 300 DPI)

Spatial projections of tallest ecotypes. Autoecology of *Pinus halepensis* distribution (left), spatial projections of tallest ecotypes for tree height (m) at age 15 years across the entire species' distribution range in the Mediterranean basin, considering maximum occupancy of each ecotype (center), and percentage of the area where each ecotype shows height superiority ($P < 0.05$) at the origin, i.e., under local occupancy (right). (a) Reference conditions (period 1961-1990 AD) based on WorldClim data (Hijmans et al., 2005). (b) Future conditions (period 2071-2100 AD) using the climate layers for RCP 4.5. (c) Future conditions using the climate layers for RCP 8.5. In left panels, trials are depicted as black triangles (up triangles: training trials; down triangles: test trials) and populations as colored squares [populations and trials represented in (b) and (c) are based on reference climate]. In central panels, the pie charts represent the frequency of each ecotype significantly outperforming ($P < 0.05$) other ecotypes for each climate condition considering maximum occupancy. The species range is derived from the EUFORGEN distribution map (48) for a resolution of 10'. See the legend of Figure 1 for ecotype acronyms.

254x190mm (300 x 300 DPI)

Spatial projections for tree height (m) at age 15 under reference conditions (period 1961-1990 AD). (a) Species level (General Performance) and (b, c, d, e, f) ecotype level, considering maximum occupancy in the Mediterranean basin. The histograms depict the class frequencies (%) of tree height at the ecotype and general level for the entire species' distribution. The class frequencies of tree height at the geographic origin of each ecotype, estimated for a 25 km radius around each ecotype's populations, are also shown in the histograms as hatched bars. A Median test (Z) for comparison between local (L) and regional (R) performance, together with their median values (M, in m), is included. Positive Z values denote local level is superior to regional level performance.

254x190mm (300 x 300 DPI)

Relative tree height differences at age 15 between future and reference conditions in the Mediterranean basin. Left panels. Maps of tree height differences (%) at the species level (General Performance, GP) for (a) RCP 4.5 scenario and (b) RCP 8.5 scenario between future (period 2071- 2100 AD) and reference conditions (period 1961-1990 AD) across the distribution range of *P. halepensis*. Right panels. Pie charts summarizing the relative importance of height differences (enhancement [$> 10\%$], inhibition [$< -10\%$], stability [-10% , $+10\%$]) at the local (left charts) and regional (i.e., maximum occupancy of each ecotype, right charts) levels for general performance and for each ecotype. See the legend of Fig. 1 for ecotype acronyms and Fig. S7 in SI for absolute tree height differences.

254x190mm (300 x 300 DPI)