

# Comparison of four European endangered red sheep based on fleece characteristics

P.-M. Parés<sup>1</sup>, R. Perezgrovas<sup>2</sup> and J. Jordana<sup>3</sup>

<sup>1</sup>*Dept. de Producció Animal, Ciència i Salut Animal, Universitat de Lleida, Lleida, Catalunya, Spain;* <sup>2</sup>*Instituto de Estudios Indígenas, Universidad Autónoma de Chiapas (IEI-UNACH), Centro Universitario Campus III, San Cristóbal de Las Casas, Chiapas, México;* <sup>3</sup>*Dep. de Ciència Animal i dels Aliments, Facultat de Veterinària, Universitat Autònoma de Barcelona, Bellaterra, Catalunya, Spain*

## Summary

Patterns of wool variations found in modern sheep breeds may shed some light on the history of circum-Mediterranean migrations, which at present are still largely unknown. In this study, the characteristics of wool from five different local European breeds were investigated. Wool samples were taken from the following sheep breeds: Aranese, Roussillon Red, Castillonnais, Levant Red and Majorcan Red. Specimens were collected from 88 animals in different herds for each breed. The following sheep fleece characteristics were investigated: length of long-coarse, short-fine and kemp (medullated) fibres; their proportion within the staple; yield after scouring with isoalcohol; and fibre diameter. The amount of short-fine fibres was lower in the Levant Red and Majorcan Red breeds, with an inversely proportional amount of kemp fibres. Clean yield with isoalcohol was about 80 percent in all these breeds. Fleece characteristics offer a glimpse of possible morphologic relationships between the Levant Red and Majorcan Red that moreover exhibit a similarity in colour and a geographical proximity. Because very little is still known about the nature of the patterns of the fleece of Atlantic West and North African sheep, the importance of the African influence on existing European Mediterranean red sheep breeds remains to be clarified. A plausible hypothesis is that different influences would have left different traces, with a lesser impact on Pyrenean breeds.

**Keywords:** *Aranese, Castillonnais, Levantina Red, Majorcan Red, Roussillon Red*

## Résumé

Les modèles des variations de la laine découverts dans les races modernes de moutons pourraient apporter quelques éclaircissements sur l'histoire des migrations autour de la Méditerranée qui sont à présent encore largement inconnues. Dans cette étude, on a examiné les caractéristiques de la laine de cinq différentes races locales européennes. On a pris des échantillons de laine des races suivantes de moutons: Aranese, Rouge du Roussillon, Castillonnaise, Levantina Red et Roja Mallorquina. Pour chaque race, on a collecté les spécimens de 88 animaux de différents troupeaux. Les caractéristiques de la toison des moutons ayant été analysées ont été les suivantes: la longueur des fibres longues et grossières, courtes et fines et de jarre (médullaires), leur proportion dans le bouquet, le rendement après le dégraissage avec l'alcool isopropylique et le diamètre des fibres. La quantité de fibres courtes et fines était inférieure dans les races Levantina Red et Roja Mallorquina, avec une quantité inversement proportionnelle de fibres de jarre. Le rendement de la laine lavée à fond avec l'alcool isopropylique était d'environ 80 pour cent dans toutes les races. Les caractéristiques de la toison donnent un aperçu des relations morphologiques possibles entre les races Levantina Red et Roja Mallorquina qui présentent en outre une similitude de couleur et une proximité géographique. En raison des connaissances très limitées sur la nature des types de toison des moutons de l'Atlantique occidentale et de l'Afrique du Nord, l'importance de l'influence africaine sur les races européennes existantes de moutons rouges de la Méditerranée n'a pas encore été précisée. Une hypothèse plausible est que les différentes influences ont laissé des traces différentes, avec un impact moindre sur les races des Pyrénées.

**Mots-clés:** *Aranese, Castillonnaise, Levantina Red, Roja Mallorquina, Rouge du Roussillon*

## Resumen

Los modelos de variación en la lana que pueden hallarse en las razas ovinas modernas pueden arrojar cierta luz sobre la historia de las migraciones circun-mediterráneas, que aún en la actualidad están poco estudiadas. En este estudio, se investigaron las características de la lana de cinco razas locales europeas diferentes. Se obtuvieron muestras de lana de las razas Aranese, Roja del Rossellón, Castellonesa, Guirra y Roja Mallorquina, agrupando un total de 88 animales de rebaños diferentes. Fueron investigadas las siguientes características del vellón: longitud de fibras largas-bastas, cortas-finas y kemp, su proporción en la mecha, rendimiento isoalcohólico, y diámetro de las fibras. Las razas Guirra y Roja Mallorquina presentaron la menor cantidad de fibras cortas-finas, y ello en proporción inversa respecto de las fibras kemp. El desengrasado con isoalcohol apareció entorno el 80% para todas las razas. Las características del vellón reflejan posibles relaciones morfológicas entre la Guirra y la Roja Mallorquina, que además exhiben una similitud en su color y proximidad geográfica. Puesto que existe poca información de los patrones del vellón de los ovinos del Atlántico Occidental y del

Norte de África, la importancia de la influencia africana sobre las razas rojas mediterráneas europeas está aún por dilucidar. Una hipótesis plausible es que diferentes influencias hayan dejado señales diferentes en esas razas, con un menor impacto en las pirenaicas.

**Palabras clave:** *Aranesa, Castellonesa, Guirra, Roja Mallorquina, Roja del Rossellón*

Submitted 17 August 2010; accepted 19 November 2010

## Introduction

Patterns of wool variations found in modern sheep breeds may shed some light on the history of circum-Mediterranean migrations, which at present are largely unknown. But there are not many reports in the literature regarding the relationships among Mediterranean sheep. For instance, Álvarez *et al.* (2004) sampled from six Spanish sheep breeds included in the ancestral Iberian Churra sheep group, Gaouar *et al.* (2005) sampled from two Algerian breeds, El Nahas *et al.* (2008) sampled only from Egyptian breeds and Ali *et al.* (2009) also sampled from Egyptian sheep. They all analysed microsatellites or structural genes.

Moreover, there is very little information about fleece characteristics for the Mediterranean breeds. The few existing studies on comparative fleece characteristics for some of these breeds have been done by Parés (2008) and Perezgrovas *et al.* (2010). Among the authors' drawbacks is that the fleece analysis can be used not only to describe single isolated breeds, but also to contribute to the study of phenotypical groups.

Taking into account rapid genetic erosion (Scherf, 2000) and the importance of animal genetic resources for the future (Notter, 1999), it may not be possible to wait for specific and detailed information on some of the less studied Mediterranean breeds and ecotypes. In this study, we concentrate on a preliminary comparative analysis of fibre and staple characteristics for the red Mediterranean breeds with the aim of establishing fibre morphologic relationships. One of our hypotheses is that differentiation between these red breeds is clearly due to more than colour pattern alone. The approach to apply conventional multivariate analysis remains valuable as these red breeds can be viewed as relatively closely related.

## Materials and methods

### Breeds studied

Samples were taken from five different local European breeds (Figure 1): Aranese (ARA), Castillonais (CAST), Levant Red (GUIR), Roussillon Red (BERB) and Majorcan Red (ROJM), the last two being funnel tailed. Except for the ARA, they yield red wool. The ARA, which is also a medium-woolled type (Parés, 2008), was taken as a

comparative group, its coat generally being white in colour (other phenotypes are allowed, but none is red).

The ARA thrives in the Aran Valley, in the Central Pyrenees. Husbandry is similar to that for the CAST, i.e. during winter each flock is pastured at the owner's farm in the valley, with feed provided in the stall, and for four months during the summer, the flocks of sheep graze normally in the sub-Alpine and Alpine grasslands. Although it is quite possible that the origin of the Aran breed may include some elements of the Merino, it is necessary to look for the presence of other stock in its formation (Parés, 2008).

The BERB (in Spain known as the Berberina) is a semifine woolled breed from the lowlands of southwestern France, especially concentrated between Narbonne and Perpignan (Babo, 2000). The 2 percent of the breed that are black are not accepted as pure BERB (Brooke and Ryder, 1977). Head, legs and belly are bare of fleece and reddish-brown (Brooke and Ryder, 1977). The origin of the breed is unknown. Pujol (1974) accepts as valid the traditional belief that the BERB is of North African origin, introduced into France via Spain. Sánchez Belda (1976a, 1976b) postulates that both the BERB and the ROJM have Barbary ancestry, and he also rejects the suggestion that the BERB and the



**Figure 1.** Origin of studied breeds: 1, ARA; 2, BERB; 3, CAST; 4, GUIR; 5, ROJM.

GUIR are closely related. Babo (2000) puts forward the possibility of Solognote or Cobourg influence. In France, the breed has been declining for many years: From more than 50 000 in 1900 (Pujol, 1974) to 3 000 in 1963 (Ministere de l'Agriculture, 1963), but its survival is not at critical levels in Spain, where many pure flocks exist, mainly in the eastern Pyrenees of Catalonia.

The CAST (formerly Saint Gironnaise and locally known as Castelhonesa or Tête Rouge), a breed (like the ARA) from the Central Pyrenees, is found in the Lez valley, southwest of St Geronç (Ariège) (Babo, 2000). The main breeding area is near the villages of Castillon-en-Couserans and Bordes (Brooke and Ryder, 1978). The Castillon breed is described by Quittet (1965), Mason (1967) and Babo (2000): Coat is white; the head, legs, and belly are reddish-brown or partly reddish-brown; the shade varies from dark to light. Lambs are born entirely reddish-brown, and mature animals are not considered pure-bred unless the head is at least partly red, a recessive characteristic (Brooke and Ryder, 1978). It has received influences from the Southdown breed (Babo, 2000). With a count of about 1 000 animals, its survival is critical (Babo, 2000).

The GUIR (also known as Guirra and Sudat) sheep is an autochthonous breed distributed throughout southeast Spain. The GUIR is characterized by a reddish coat colour and very oily wool (Sánchez Belda, 1976a, 1976b; Esteban, 2003) ("Guirra" is the word for "reddish" in the Valencian dialect; "Sudat" means "greasy", in reference to the oily condition of the uncleaned wool – Sánchez Belda, 1976a, 1976b; Brooke and Ryder, 1977). Its numbers have been decreasing in recent years (Esteban, 2003). The colour of the fleece of the animals varies from reddish-brown to yellow-white (Sánchez Belda, 1976a, 1976b; Brooke and Ryder, 1977). The colour of newborn lambs is dark reddish-brown (Sánchez Belda, 1976a, 1976b; Esteban, 2003). With age, the coat lightens, becoming a dirty cream colour at full maturity (Sánchez Belda, 1976a, 1976b; Brooke and Ryder, 1977). Both sexes lack horns (Sánchez Belda, 1976a, 1976b). Years ago it was believed that the GUIR breed was a pigmented variety of the Segureña breed or a similar breed to the Manchega variety (Sánchez Belda, 1976a, 1976b), but nowadays it is considered ethnically independent with African ancestry, consistent with the close ties between the Spanish central Mediterranean coast and the Maghreb (Sánchez Belda, 1976a, 1976b). Specifically, it would have originated on the Atlantic coast of Morocco from the Beni Ahsen breed, which has exactly the same colour and wool characteristics (Sánchez Belda, 1976a, 1976b; Esteban, 2003).

The ROJM (also known as Coete) is a breed found on the island of Majorca. In recent decades, the breed has steadily declined to such an extent that its survival is uncertain (Esteban, 2003). The ROJM developed from crossing the island animals with funnel-tailed and chestnut North African breeds (Sánchez Belda, 1976a, 1976b; Esteban,

2003). Both sexes lack horns (Sánchez Belda, 1976a, 1976b). The tail is funnel shaped and the wool has hairy fibres (Esteban, 2003).

Additional breed descriptions are available from Esteban (2003) for Spanish breeds and Babo (2000) for French ones.

Meat and/or milk production are the unique traits for these sheep, with wool production being unimportant. Farmers shear their flocks once a year between late April and June.

## Sampling

Adult animals were randomly chosen (a herd for each breed). Staples were taken before shearing from the shoulder and stored in plastic bags for further analysis. Subsamples were taken to measure fibre length and diameter. A total of 88 animals were sampled.

To identify differences between fibres, macroscopic and microscopic features were analysed using the method as described in Rojas *et al.* (2005).

## Traits studied

The following traits were considered in the study: length of long-coarse, short-fine and kemp fibres; percentage of long-coarse, short-fine and kemp fibres (staple composition); percentage of clean yield using isoalcohol; fibre:kemp ratio; average fibre diameter; and F30 (fibres with diameter >30 µm). Formal statistical requirements, such as covariance homogeneity between samples and multivariate normality of distribution, are assumed to be met.

A more precise view of relationships can be gleaned from the similarity coefficients. So to compare differences between breeds, a principal coordinate (PCO) scatter diagram and a paired cluster analysis were done based on the Gower distance matrix, considering each breed as a group. F30 was not considered in this case. Used data from ARA, CAST and BERB were previously compiled by Parés (2008) and data from GUIR and ROJM by Perezgrovas *et al.* (2010). All crude values are available from the first author upon request. Data were analysed using PAST<sup>®</sup> v. 1.94b ([http://palaeo-electronica.org/2001\\_1/past/issue1\\_01.html](http://palaeo-electronica.org/2001_1/past/issue1_01.html)).

## Results and discussion

The descriptive statistics obtained are shown in Tables 1 and 2. Statistical differences for some of the studied traits were detected. The results for the breeds studied showed an overall fibre diameter of  $29.5 \pm 10.65$  µm. The rather symmetrical diameter distribution makes them a short, fine wool type. The histograms for frequency of diameter distribution (not presented here) confirm the macroscopic findings, with a lower proportion of long-coarse fibres, although ROJM also shows the highest percentage of long-coarse fibres, indicating a slight tendency towards the

**Table 1.** Descriptive statistics obtained from fleece analysis (mean  $\pm$  SE) (breed abbreviations in the text).

	ARA	BERB	CAST	GUIR	ROJM
No. of samples	18	15	28	15	12
Long-coarse fibres length	4.17 $\pm$ 0.43 <sup>a</sup>	2.56 $\pm$ 1.40 <sup>b</sup>	4.33 $\pm$ 1.75 <sup>c</sup>	7.55 $\pm$ 0.75 <sup>d</sup>	12.04 $\pm$ 2.2 <sup>e</sup>
Short-fine fibres length	4.10 $\pm$ 1.12 <sup>a</sup>	2.94 $\pm$ 0.55 <sup>b</sup>	4.64 $\pm$ 1.06 <sup>c</sup>	6.30 $\pm$ 0.95 <sup>ac</sup>	9.54 $\pm$ 1.48 <sup>d</sup>
Kemp length	1.28 $\pm$ 0.49 <sup>a</sup>	0.88 $\pm$ 0.28 <sup>B</sup>	0.85 $\pm$ 0.46 <sup>c</sup>	2.40 $\pm$ 1.00 <sup>d</sup>	2.92 $\pm$ 0.97 <sup>d</sup>
% long coarse	5.85 $\pm$ 13.91 <sup>ab</sup>	0.75 $\pm$ 2.40 <sup>b</sup>	1.78 $\pm$ 6.20 <sup>b</sup>	3.94 $\pm$ 3.50 <sup>ab</sup>	10.33 $\pm$ 5.12 <sup>a</sup>
% short fine	93.17 $\pm$ 13.97 <sup>a</sup>	98.35 $\pm$ 2.60 <sup>a</sup>	97.01 $\pm$ 6.07 <sup>a</sup>	92.36 $\pm$ 4.85 <sup>a</sup>	83.42 $\pm$ 7.95 <sup>b</sup>
% kemp	0.96 $\pm$ 1.33 <sup>a</sup>	0.88 $\pm$ 0.85 <sup>a</sup>	1.19 $\pm$ 1.05 <sup>a</sup>	3.69 $\pm$ 2.65 <sup>b</sup>	6.23 $\pm$ 5.00 <sup>c</sup>
Isoalcoholic yielding (%)	84.29 $\pm$ 9.48 <sup>ac</sup>	81.19 $\pm$ 10.06 <sup>bc</sup>	80.59 $\pm$ 7.71 <sup>bc</sup>	74.71 $\pm$ 5.92 <sup>b</sup>	76.50 $\pm$ 5.88 <sup>bc</sup>
Fibres:kemp	102.30 $\pm$ 17.01 <sup>ab</sup>	121.06 $\pm$ 27.13 <sup>a</sup>	117.67 $\pm$ 27.30 <sup>a</sup>	42.62 $\pm$ 9.44 <sup>ab</sup>	30.54 $\pm$ 7.93 <sup>b</sup>

Fibre lengths in cm.

Different letters in the same row indicate significant differences ANOVA ( $P < 0.05$ ) among each breed.

double coat (important percentages of both long-coarse and short-fine fibres). It is also surprising that the ROJM, with the highest percentages of long-coarse and kemp fibres, has a more uniform fleece in terms of SD of fibre diameter than the CAST and GUIR and a more desirable value for F30 than the CAST. All can be classified as “entrefine” wool (medium quality, fibre diameter from 24 to 36  $\mu\text{m}$ ) according to the Spanish classification, with the CAST wool being the coarsest (as described by Quittet, 1965; Mason, 1967) and with kemp, and ARA being the finest with nearly no kemp. GUIR and ROJM show the lowest yield after clean yielding with isoalcohol (74.7 and 76.5percent, respectively), thus confirming the “oily wool” that characterizes these breeds (Perezgrovas *et al.*, 2010). The fleece of GUIR is also the least uniform among the studied breeds in fibre diameter. Its fibre lengths appeared slightly longer than those reported by Esteban (2003), which are of 5–6 cm. Obtained fibre lengths for ROJM appeared too longer than those reported by Esteban (2003), which are of 8–9 cm.

Table 3 shows the Gower distances between breeds. Figure 2 presents the PCO scatter diagram. The results of the PCO analysis are eminently interpretable in terms of function of

all analysed traits. The first PCO accounts for about three-quarters of the total variance. The first component variate explains the majority of the variance (79.5percent) and acts to separate the two Iberian red breeds (GUIR and ROJM) from the French ones (CAST and BERB).

Figure 3 displays the paired clustering obtained ( $\rho = 0.941$ ). Iberian and French breeds are widely separated from each other, GUIR and ROJM having a low fibre-to-kemp ratio (high percentage of kemp and low percentage of short-fine fibres). Their fibre diameter also tends to be lower; the wool is markedly greasy and the staple longer. GUIR and ROJM could be grouped as “hairy and high-greasy medium-wool” breeds. BERB and CAST could be defined as “low-greasy medium-wool” breeds with the CAST having very coarse fibres.

The results of the clustering analysis here are similar to one of those done by Parés (2008) with ARA and French breeds, but differ slightly from Parés (2010) who investigated 31 Iberoamerican breeds (that include the studied breeds in this research). One reason that the results of this study differ is that the resulting dendrogram may strongly vary according to the selected clustering methods.

**Table 2.** Some descriptive traits obtained from fibre diameter analysis (breed abbreviations in the text).

	ARA	BERB	CAST	GUIR	ROJM
No. of fibres	2 184	1 248	1 380	1 560	1 248
Min ( $\mu\text{m}$ )	9	10	11	12	14
Max ( $\mu\text{m}$ )	135	93	145	175	80
Mean ( $\mu\text{m}$ )	29.5 <sup>b</sup>	26.7 <sup>a</sup>	35.3 <sup>c</sup>	27.7 <sup>ac</sup>	27.9 <sup>dc</sup>
Std. error	0.175	0.175	0.287	0.372	0.267
Variance	67.56	38.50	113.71	216.40	89.43
Stand. dev	8.22	6.21	10.66	14.71	9.46
CV (%)	27.77	23.21	30.18	53.04	33.82
Median ( $\mu\text{m}$ )	29	26	34	25	26
25 percentile	24.00	23.00	28.00	22.00	22.00
75 percentile	34.00	30.00	41.00	29.00	31.75
Skewness	2.734	2.178	2.014	5.294	2.078
Kurtosis	154.421	19.355	-29.254	-287.537	6.196
Geom. mean	28.58	26.07	33.91	25.91	26.72
F30	39.42	23.48	66.81	19.49	27.40

All fibre types are included.

Different letters in the row indicate significant differences ANOVA ( $P < 0.05$ ) between means among each breed.


**Table 3.** Matrix of Gower distances (abbreviations in the text).

	ARA	BERB	CAST	GUIR	ROJM
ARA	0.000				
BERB	0.254	0.000			
CAST	0.251	0.201	0.000		
GUIR	0.429	0.521	0.518	0.000	
ROJM	0.704	0.847	0.838	0.368	0.000

**Discussion**

The very early breeds of sheep had two coats: an outer coat of kemp and an under coat of fine wool or down that is shed in the spring (McManus, Rezende and Olegário de Araújo, 2010). The kemp has been gradually bred out of most sheep so that they have coats that are all wool (McManus, Rezende and Olegário de Araújo, 2010). Fleece characteristics offer a glimpse of possible historic relationships among the breeds studied beyond their similarity in colour and geographical proximity.


The geographic location of the Iberian Peninsula makes it a natural link between Europe and North Africa. However, it is a matter of debate to what extent African influences via the Strait of Gibraltar affected Iberia’s prehistoric development. Because early African pastoralist communities were dedicated to sheep breeding, a possible means of detecting prehistoric African–Iberian contacts might be to analyse the origin of ovine breeds on the Iberian Peninsula. But because we still know very little about the nature of the fleece patterns of Atlantic West and North African breeds,



**Figure 3.** Paired cluster analysis based on the Gower distance matrix depicting the results of the cluster analysis described in the text. Each breed is considered as a group (breed abbreviations in the text). Note the initial bifurcation between Iberian red breeds (GUIR and ROJM) and French ones (CAST and BERB). Note also that GUIR and ROJM are in a cluster of long branches.  $\rho = 0.941$ .

the importance of the African influence on modern European red sheep remains to be clarified.

Evidence from the Cairo Genizah indicates quite clearly that in the eleventh and twelfth centuries the Mediterranean world was a kind of medieval common



**Figure 2.** PCO scatter diagram of the first two PCOs based on the Gower distance matrix, considering each breed as a group (breed abbreviations in the text). Note that on the 1 coordinate (explaining the majority of the total variance) the two Iberian red breeds (GUIR and ROJM) are very far separated from the French ones (CAST and BERB) and ARA.

market, forming a free trade area (Goitein, 1967). This communications network, shared by Christians, Jews and Moslems, expressed the notion that there was blessing in movement “fi'l-haraka baraka” (Glick, 1979). In addition, the Atlantic maritime trade between Spain, Portugal and the Maghreb at this time is well documented by Picard (1997). All these influences would have left different traces, with a less frequent influence on Pyrenean breeds. It seems possible that different breeds, as they spread northward, were the origin of different European breeds. Subsequent selective breeding for desirable traits such as wool, milk and meat production as well as environmental tolerance was responsible for the development of more differentiated sheep populations. A loss of wool improvement can be assumed, at least in recent decades. The influx of stocks through different pathways is compatible with this hypothesis as the southernmost breeds (GUIR and ROJM) displaying a high degree of interbreed Gower distances. Moreover, it seems difficult to suppose that the funnel-tailed ROJM and the thin-tailed GUIR, which appear in the same cluster, are closely related. Because very little is still known about the nature of the patterns of the fleece of Atlantic West and North African sheep, the importance of the African influence on existing European red sheep breeds remains to be clarified. A plausible hypothesis is that different African influences would have left different traces, with a lesser impact on Pyrenean breeds. DNAmT finds could further clarify the links between African and Iberian cultures.

More red sheep breeds from the Mediterranean basin and Atlantic African coast had to be included in further studies to achieve a good general study of red sheep breeds: Beni Ahsen breed (Atlantic Morocco); Anatolian Red and Red Karaman breeds (Turkey); Tunisia, Libya, Lebanon and Algeria sheep, etc.

## Conclusions

A shortcoming of the methodology used here is that the sampling scheme did not include all the red sheep breeds in the geographical area under study (Mediterranean basin). Nevertheless, the characterization of wool in these five sheep breeds will allow comparison with other breeds as done by Parés (2008) and Parés (2010).

Other traits as staple colour, as used by Perezgrovas *et al.* (2010), would be tested as discriminative parameters in view of comparing breeds.

## Acknowledgements

Very special thanks are extended to the Spanish and French farmers who allowed us to collect samples from their herds. Special thanks are offered to the staff at CEAGA in Castielfabib (Rincón de Ademuz, Valencia) for providing the superior facilities for sampling their

GUIR animals and to Guillem Riera from the Universitat de Lleida, who kindly supplied us with ROJM samples.

## References

- Ali, B.A., El-Hanafy, A.A. & Salem, H.H. 2009. Genetic biodiversity studies on IGFBP-3 gene in Egyptian sheep breeds. *Biotechnol. Anim. Husbandry*, 25(1–2): 101–109.
- Álvarez, I., Royo, L.J., Fernández, I., Gutiérrez, J.P., Gómez, E. & Goyache, F. 2004. Genetic relationships and admixture among sheep breeds from Northern Spain assessed using microsatellites. *J. Anim. Sci.*, 82: 2246–2252.
- Babo, D. 2000. *Races ovines et caprines françaises*. Edits. Paris, France Agricole.
- Brooke, C.H. & Ryder, M.L. 1977. Characteristics of some less-common breeds of sheep in Southern Europe: a preliminary survey. *Ann. Génét. Sél. Anim.*, 9(2): 163–180.
- Brooke, C.H. & Ryder, M.L. 1978. Declining breeds of mediterranean sheep. *FAO Animal Production and Health Paper* 8. Rome.
- El Nahas, S.M., Hassan, A.A., Mossallam, A.A.A., Mahfouz, E.R., Bibars, M.A., Oraby, H.A.S. & de Hondt, H.A. 2008. Analysis of genetic variation in different sheep breeds using microsatellites. *Afr. J. Biotechnol.*, 7(8): 1060–1068.
- Esteban, C. 2003. *Razas ganaderas españolas. II. Ovinas*. Madrid, Ministerio de Agricultura, Pesca y Alimentación.
- Gaouar, S., Tabet-Aoul, N., Derrar, A., Goudarzy-Moazami, K. & Saïdi-Mehtar, N. 2005. Genetic diversity in Algerian sheep breeds, using microsatellite markers. In H.P.S. Makkar & G.J. Viljoen, eds. *Applications of gene-based technologies for improving animal production and health in developing countries*. The Netherlands, Springer. pp. 641–644.
- Goitein, S.D. 1967. *A Mediterranean society*. Vol. I. *Economic foundations*. Los Angeles, CA, University of California Press.
- Glick, T.F. 1979. *Islamic and Christian Spain in the early middle ages*. Princeton, NJ, Princeton University Press.
- McManus, C., Rezende, S. & Olegário de Araújo, R. 2010. Genetics and breeding of sheep in Brazil. *R. Bras. Zootec.* 39(supl. especial): 236–246.
- Mason, I.L. 1967. *The sheep breeds of the Mediterranean*. Farnham Royal, England, FAO and Commonwealth Agricultural Bureaux.
- Ministere de l'Agriculture. 1963. *Le Cheptel Bovin, Porcin*. Paris, Ovin et les Productions.
- Notter, D.R. 1999. The importance of genetic diversity in livestock populations of the future. *J. Anim. Sci.*, 77(1): 61–69.
- Parés, P.M. 2008. *Caracterització estructural i racial de la raça ovina Aranese*. Univ. Autònoma de Barcelona, Spain. (PhD dissertation)
- Parés, P.M., Perezgrovas, R., Jordana, J. 2010. Diversity and breed comparison of wool parameters in 31 different American and European ovine breeds. In *5th European Symposium in South American Camelids and First European Meeting on Fibre Animals*, Sevilla (Spain), 2010 (in press).
- Perezgrovas, R., Parés, P.M., Rodríguez, G. & Zaragoza, L. 2010. *Características de la mecha y la fibra de lana en ovejas Guirra y Roja Mallorquina de España*. *Simp. Iberoamericano sobre Conservación y Utilización de Recursos Zoogenéticos*. Paraíba (Brazil) (in press).
- Picard, C. 1997. *L'océan Atlantique musulman; De la conquête arabe à l'époque almohade; Navigation et mise en valeur des côtes d'al-Andalus et du Maghreb occidental (Portugal-Espagne-Maroc)*. Paris, Maisonneuve et Larose.

- Pujol, T.P.** 1974. *La Race Rouge du Rousillon*. Paris, Institut Technique de l'Élevage Ovine et Caprin (ITOVIC).
- Quittet, E.** 1965. *Races Ovines Françaises*. Paris, Maison Rustique.
- Rojas, A.L., Perezgrovas, R., Rodríguez, G., Russo-Almeida, P. & Anzola, H.** 2005. Caracterización macro y microscópica de la lana en ovinos autóctonos iberoamericanos de vellón blanco. *Arch. Zootec.*, 54: 477–483.
- Sánchez Belda, A.** 1976a. Los ovinos de raza roja levantina. *Av. Aliment. Mejora Anim.*, 4: 3–11.
- Sánchez Belda, A.** 1976b. Los ovinos de raza Roja Mallorquina de cola ancha. *Av. Aliment. Mejora Anim.*, 1: 3–9.
- Scherf, B.D.** 2000. *World watch list for domestic animal diversity*. 3rd ed. Rome, FAO.