

**Universitat de Lleida**

Document downloaded from:

<http://hdl.handle.net/10459.1/66422>

The final publication is available at:

<https://doi.org/10.1007/s11947-018-2217-z>

Copyright

(c) Springer Science+Business Media, LLC, part of Springer Nature, 2018

# Food and Bioprocess Technology

## Effect of ultrasound pre-treatment on the physical, microbiological, and antioxidant properties of calçots

--Manuscript Draft--

<b>Manuscript Number:</b>	FABT-D-18-00210R2	
<b>Full Title:</b>	Effect of ultrasound pre-treatment on the physical, microbiological, and antioxidant properties of calçots	
<b>Article Type:</b>	Original Research	
<b>Keywords:</b>	Allium cepa L.; thermal processing; Gastrointestinal digestion; antioxidant activity; novel technologies	
<b>Corresponding Author:</b>	Ingrid Aguiló-Aguayo, PhD IRTA Lleida, SPAIN	
<b>Order of Authors:</b>	Lorena Zudaire Tomás Lafarga Inmaculada Viñas Maribel Abadías Nigel Brunton Ingrid Aguiló-Aguayo, PhD	
<b>Corresponding Author Secondary Information:</b>		
<b>Corresponding Author's Institution:</b>	IRTA	
<b>Corresponding Author's Secondary Institution:</b>		
<b>First Author:</b>	Lorena Zudaire	
<b>First Author Secondary Information:</b>		
<b>Order of Authors Secondary Information:</b>		
<b>Funding Information:</b>	Agència de Gestió d'Ajuts Universitaris i de Recerca (FI-2017-B2-00164) Secretaría de Estado de Investigación, Desarrollo e Innovación (FJCI-2016-29541) Secretaría de Estado de Investigación, Desarrollo e Innovación (RYC-2016-19949)	Dr. Lorena Zudaire Dr. Tomás Lafarga Dr. Ingrid Aguiló-Aguayo
<b>Abstract:</b>	<p>The effect of ultrasound (US) treatment (40 kHz, 250 W) for 0, 10, 25 and 45 min on the physical and microbiological quality, total antioxidant capacity (TAC) and total phenolic content (TPC) of calçots (<i>Allium cepa</i> L.) was evaluated. Moreover, the effect of roasting (270 °C, 8 min) and in vitro simulated digestion on the antioxidant properties was studied. Overall, US treatment had no effect on the physical quality and antioxidant properties of calçots regardless the treatment time, while thermal processing produced an increase on the TAC and maintenance in TPC. Furthermore, the digestion process caused a remarkable decrease on the TAC and TPC, but that decrease was higher in roasted than in fresh samples. The microbial load of all US-treated fresh samples was below 6 log (cfu g<sup>-1</sup>) and a decrease of 1-log reduction was observed after treating for 45 min. Those results indicated that US pre-treatment had no negative effects on the quality of calçot while produced a decrease on the microbial load at high processing times.</p>	

Firstly, all issues mentioned in the reviewer's comments has been considered. All changes has been marked in yellow colour. Manuscript, tables and figures have been revised taking into account reviewer's comments. Then, comments from the editors and reviewers are explained (new paper's lines):

**Reviewer #6:**

**1. Ultrasound is widely used in food industry. I suggest the author evaluate the economic cost. Because we are not sure if US treatment will increase production efficiency and decrease the cost of production. Therefore, we are not sure if US treatment deserves to be popularized and be used in food production.**

Recently, Welti-Chanes et al. (2017) have published a review where they remarked that 'more than 1000 scientific publications indicate that this treatment offers a net advantage in terms of productivity, yield, and selectivity'. Moreover, Gallo et al. (2018) highlighted that 'competitive energy costs and low maintenance make ultrasound processed economically profitable'.

We added 'only consumed a fraction...Wang et al. 2015; Welti-Chanes et al. 2017)' (Lines 39-43, page 2) emphasising the industrial and economical advantages of ultrasound application in food industry.

**2. In 2.2 sonication, the author said "The surface of water (tap water) in the bath was kept at the same level during each experiment but without temperature controller". As I know, under the US treatment of 250W, 45min, the water will be much hotter than T0. Does the temperature of water induced by different treatment time affect the detection indicators?**

We measured the temperature before and after of each treatment, but we decided not to put in the manuscript because there were no significant differences in most of the studied parameters. We considered an irrelevant information for the interpretation and discussion of the results.

Time (min)	$\Delta T$ (°C)
10	$2.67 \pm 0.23$
25	$5.33 \pm 0.50$
45	$10.40 \pm 1.97$

$$\Delta T = \text{Post-US Temperature (}^{\circ}\text{C)} - \text{Pre-US Temperature (}^{\circ}\text{C)}$$

**3. The author said that US treatment has some effects on physical properties of calcots. I suggest that the author could make paraffin section to show the tissue structure of calcots. It would be another way to show the effects of US treatment on calcots physical properties.**

We agree with that comment, but we did not take a photograph of transversal section of any *calçot*. We will consider this recommendation for future projects and publications. We only took a photo before and after each treatment, but they were photos of general appearance. For example:

- **Before** Ultrasound treatment:



- **After** Ultrasound treatment:



10 min

25 min

45 min

**4. The Browning Index (BI) only appeared in the Table 1. The author didn't mention it in the discussion texts. If the author put Browning Index in results, it is necessary to give comment on it.**

We agree with that comment. We added 'Moreover, BI values...among them' (Lines 160-161, page 8).

**5. Line 106: the author mentioned the dry matter (DM), but this abbreviation was used only one time in the following sentence. In the following texts the author used another notion dry weight (dw). It would be better to unify the abbreviation.**

We agree with that comment. We decided to put dry weight (dw) throughout the document. We put 'weight (dw)' instead of 'matter (DM)' in line 109 of page 5.

## References

- Gallo, M., Ferrara, L., & Naviglio, D. (2018). Application of Ultrasound in Food Science and Technology: A Perspective. *Foods*, 7(10), 1–18. doi:10.3390/foods7100164
- Welti-Chanes, J., Morales-de la Peña, M., Jacobo-Velázquez, D. A., & Martín-Belloso, O. (2017). *Opportunities and Challenges of Ultrasound for Food Processing: An Industry Point of View*. *Ultrasound: Advances for Food Processing and Preservation*. Academic Press. doi:10.1016/B978-0-12-804581-7.00019-1

[Click here to view linked References](#)

1           **Effect of ultrasound pre-treatment on the physical, microbiological, and**  
2           **antioxidant properties of calçots**

3           Lorena Zudaire<sup>1</sup>, Tomás Lafarga<sup>1</sup>, Inmaculada Viñas<sup>2</sup>, Maribel Abadías<sup>1</sup>, Nigel  
4           Brunton<sup>3</sup>, Ingrid Aguiló-Aguayo<sup>1\*</sup>

5           <sup>1</sup> IRTA, XaRTA-Postharvest, Parc Científic i Tecnològic Agroalimentari de Lleida, Parc  
6           de Gardeny, Edifici Fruitcentre, 25003, Lleida, Catalonia, Spain.

7           <sup>2</sup> Food Technology Department, University of Lleida, XaRTA-Postharvest, Agrotecnio  
8           Center, Lleida, Spain

9           <sup>3</sup> University College Dublin, School of Agriculture and Food Science, Dublin 4, Ireland

10          \* Dr Ingrid Aguiló-Aguayo: [ingrid.aguilo@irta.cat](mailto:ingrid.aguilo@irta.cat), (+34) 902 789 449, Ext.: 1551

11          ORCID:

- 12          • Lorena Zudaire: 0000-0001-9761-1155
- 13          • Tomas Lafarga: 0000-0002-1923-7214
- 14          • Inmaculada Viñas: 0000-0001-5182-2520
- 15          • Maribel Abadías: 0000-0003-0113-8979
- 16          • Nigel Brunton: 0000-0002-5893-1751
- 17          • Ingrid Aguiló-Aguayo: 0000-0002-4867-1554

19          **Abbreviations**

20           $\Delta E^*$ : Colour difference; BI: Browning Index; DPPH: 2,2-diphenyl-1-picrylhydrazyl; FRAP:  
21          Ferric Reducing Antioxidant Power;  $h^\circ$ : Hue angle; TAC: Total Antioxidant Capacity; TPC: Total  
22          Phenolic Content; US: Ultrasound.

## 23 1. Introduction

24 *Calçots* (*Allium cepa* L.) are the immature floral stems of second-year onion resprouts of the  
25 ‘Ceba Blanca Tardana de Lleida’ onion landrace. The singularity of the production of this product  
26 has helped to confer protected status from the European Union and ‘Calçot de Valls’ being  
27 awarded with the Protected Geographical Indication (EC No 905/2002) (Simó et al. 2013; Zudaire  
28 et al. 2017). An increased demand and interest for *calçots* has motivated researches to explore  
29 new postharvest techniques such as minimal processing or ultrasound (US) treatment, thus  
30 maintaining their physical, microbiological, and nutritional quality.

31 Thermal pasteurization and sterilization are two common techniques used for the inactivation of  
32 microorganisms in food products. However, the effectiveness of those methods is based on long  
33 exposure time and high temperatures, which generally results in a deterioration in functional  
34 properties, sensory characteristics, and nutritional value (Piyasena et al. 2003). In recent years,  
35 emerging non-thermal technologies, such as high pressure, pulsed electric fields, ultraviolet light,  
36 intense pulsed light, and US treatments, have been widely studied for application in food industry  
37 (de São José et al. 2014). High energy (high power, high-intensity) US are usually applied in the  
38 food industry with frequencies ranging between 20 and 100 kHz. This technology has become an  
39 attractive option for food processors because **only consumed a fraction of the time and energy**  
40 **normally need for traditional processes, reduces processing cost, guarantees food safety, improves**  
41 **food quality, reduced chemical and physical risks, and is considered environmentally friendly** ~~the~~  
42 ~~acoustic waves are generally considered safe, non-toxic, and environmentally friendly~~ (Awad et  
43 ~~al. 2012; Chemat et al. 2011; Wang et al. 2015; Welte-Chanes et al. 2017).~~

44 Previous studies suggested US processing as a promising technology if it used as an auxiliary pre-  
45 treatment to sanitizers in reducing initial microbial populations of foods (Ding et al. 2015).  
46 However, the effect of US on the total antioxidant capacity (TAC) of food is a controversial issue.  
47 On the one hand, the generation of reactive oxygen species such as hydroxyl radicals could affect  
48 the quality of some foods by reducing the TAC (Kentish and Ashokkumar 2011). On the other

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49 hand, those species could impose oxidative stress to fresh products and hence, induce the TAC of  
50 fruits and vegetables (de São José et al. 2014). For example, the application of US (20 kHz, 400  
51 W) for 10 min had no remarkable effect on the TAC and total phenolic content (TPC) of  
52 mushrooms (Lagnika et al. 2013). However, TPC of minimally processed pineapples increased  
53 after US treatment (37 kHz) at 25 or 29 W for 10-15 min (Yeoh and Ali 2017).

54 Many vegetables including *calçots*, onion, or carrots can be either eaten raw or after cooking.  
55 *Calçot* are usually eaten after roasting process. Culinary processes produce significant changes  
56 such as degradation of thermolabile compounds and formation others due to heat-induced  
57 chemical reactions. Roasting could affect phenolic compounds and, consequently the TAC of  
58 foods (Juániz et al. 2016). Furthermore, the TPC and TAC of fruit and vegetables could also be  
59 affected by the human digestion process. During gastrointestinal digestion, polyphenols could  
60 suffer changes due to their interaction with other food components, degradation or metabolization.  
61 These structural changes could affect both their uptake and bioactivity and hence, the TAC  
62 (Bouayed et al. 2012).

63 The objective of this study was to evaluate the effects of US processing for either 10, 25, or 45  
64 min on the physical and microbiological quality, TPC, and TAC of raw and roasted *calçots*.  
65 Moreover, an *in vitro* simulated gastrointestinal digestion of both raw and roasted samples was  
66 carried out to evaluate the resistance of the TAC and TPC to gastrointestinal digestion.



## 67 2. Material and Methods

### 68 2.1 Plant Material

69 *Calçots* were provided by the ‘Cooperativa Agrícola Valls’ (Tarragona, Spain) at commercial  
70 size. The *calçots* had the European quality label PGI ‘Calçot de Valls’ establishing that their  
71 diameter and size are within the legal ranges (D.A.R.P. 2009). Samples cultivated in northeast  
72 Spain (41°13’47’’N, 01°13’12’’E) during the crop growing seasons of 2016 and 2017. Pre-  
73 conditioning was conducted according to the study of Aguiló-Aguayo et al. (2015) which  
74 consisted of cutting roots and external leaves from the edible part as well as removing the outer  
75 peel. Fresh *calçots* were immersed in a 10 L bath which contained 100 mg L<sup>-1</sup> of sodium  
76 hypochlorite at room temperature under continuous agitation for 60 s. Samples were further rinsed  
77 with tap water for 1 min, dried at room temperature, and labelled as Control.

### 78 2.2 Sonication

79 Eight *calçots* for each time and repetition were directly immersed in a sonicator bath (Frequency  
80 40 kHz, Power 250 W, JP SELECTA S.A., Barcelona, Spain) and the treatment time (0, 10, 25,  
81 45 min) was varied for each batch. The surface of water (tap water) in the bath was kept at the  
82 same level during each experiment but without temperature controller (initial temperature 17 ± 1  
83 °C). All samples were weighed before and after US treatment. All samples were dried at room  
84 temperature. On each treatment time and repetition half of fresh-cut *calçots* were taken to  
85 firmness, colour and total aerobic count measurements. The rest were roasted as Zudaire et al.  
86 (2017) described. Briefly, *calçots* were roasted at 270 °C for 8 min using a Self Cooking Center  
87 (Mod SCC WE 101, Rational AG, Landsberg am Lech, Germany) and then, cooled into a blast  
88 chiller (Infrico, Cordoba, Spain) until they reached 3 °C. After conducting those assays, both fresh  
89 and roasted samples were crushed, powered and frozen with liquid nitrogen and stored at -80 °C  
90 for nutritional analysis and gastrointestinal digestion.

91

## 92 **2.3 Colour**

93 The colour of the white shaft was measured with a CR-200 Minolta Chroma Meter (Minolta,  
94 INC., Tokyo, Japan). Colour was measured using CIE L\*, a\*, b\* coordinates with illuminant D65  
95 which approximates to daylight and 10° observer angle. L\* defines the lightness, and a\* and b\*  
96 define the red-greenness and blue-yellowness, respectively. These values were used to calculate  
97 the browning index (BI) and hue angle ( $h^\circ$ ) as previously described by Liu et al. (2016) and  
98 Colás-Medà et al., (2016), respectively. Furthermore, difference from the control ( $\Delta E^*$ ) was  
99 calculated following the methodology described by Wibowo et al. (2015).

## 100 **2.4 Firmness**

101 To assess changes on texture, firmness (N) was measured at 5 cm from the roots set in transversal  
102 position using the TA.TX2 Texture Analyzer (Stable Micro Systems Ltd., Surrey, England)  
103 attached with a Warner-Blatzler blade (HDP/BSK: Blade set with knife). The sample was placed  
104 into the press holder, and then the blade was moved downwards at different rates: pre-test rate: 5  
105 mm s<sup>-1</sup>; test rate: 1 mm s<sup>-1</sup>; post-test rate: 10 mm s<sup>-1</sup> to 60 mm below the bottom of the holder.  
106 Data acquisition rate was 200 pulses per sec.

## 107 **2.5 Dry matter determination**

108 Due to differences in water content between fresh and roasted samples, total antioxidant capacity  
109 and total phenolic content calculations were made on a dry **weight (dw) matter (DM)** basis. For  
110 determination of DM content, 4-5 g of fresh or roasted sample (as triplicate) were dried in a  
111 convection oven at 105 °C for at least 40 h until reaching a constant weight.

## 112 **2.6 Determination of Total Antioxidant Capacity**

113 TAC was determined using two different methods: 2,2-diphenyl-1-picrylhydrazyl (DPPH\*)  
114 radical scavenging assay and ferric reducing antioxidant power (FRAP) assay. The extraction and

115 assays were carried out according to the methods described by Plaza et al. (2016). Results were  
116 expressed on a dry weight (dw) basis as mol of ascorbic acid equivalents per kg.

## 117 **2.7 Determination of TPC**

118 The extraction and determination of TPC were determined by the Folin-Ciocalteu method  
119 (Singleton et al. 1999), following the modifications described by Altisent et al. (2014). Results  
120 were expressed on a dry weight (dw) basis as g of gallic acid equivalent per kg.

## 121 **2.8 Microbial quality**

122 The total aerobic count of *calçots* was analysed in triplicate as described by Alegre et al. (2011).  
123 Briefly, the edible part of two *calçots* per treatment were cut and 10 g of were diluted in 90 mL  
124 of buffered peptone water (Oxoid LTD, Basingstoke, Hampshire, England) in a sterile bag and  
125 homogenized in a masticator paddle blender (IUL Masticator Basic 400 ml, IUL Instruments,  
126 Barcelona, Spain) at 250 impact s<sup>-1</sup> for 90 s in triplicate. Further ten-fold dilutions were made  
127 with saline peptone (SP; 8.5 g L<sup>-1</sup> NaCl and 1 g L<sup>-1</sup> peptone). Aliquots of serial dilutions were  
128 spread in duplicate onto plates with Plate Count Agar (Biokar Diagnostics, Beauvais, France) and  
129 were incubated at 30 ± 1 °C for 3 d. The results were represented as log colony forming units  
130 (cfu) per gram basis on fresh weight. Microbiological analyses were performed in triplicate.

## 131 **2.9 In vitro gastrointestinal digestion**

132 *In vitro* gastrointestinal digestion was performed according to the method described by Minekus  
133 et al. (2014) with minors modifications (Zudaire et al. 2017). The simulated digestion was  
134 performed in triplicate for each treatment for raw and roasted samples. A blank was prepared  
135 using only distilled water instead of sample following the same procedure. Results were compared  
136 with non-digested samples. Determinations of TAC using both the FRAP and DPPH<sup>•</sup> methods  
137 and TPC were performed after digestion.

138

139 **2.10 Statistical analysis**

1  
2  
3 140 All data were firstly evaluated for normal distribution (Shapiro-Wilk W Test) and homogeneity  
4  
5 141 of variance (Levene's Test) of residues. Significant differences between results were calculated  
6  
7 142 by using one-way analysis of variance (ANOVA). In case of non-normality or unequal variances  
8  
9 143 the non-parametric equivalents (Wilcoxon/Kruskal-Wallis Tests) were used. Differences were  
10  
11 144 ~~considered to be~~ were significant at  $p < 0.05$  (95 % confidence level). In case of significant  
12  
13 145 differences, multiple comparison of means was established with the Post Hoc Tukey-Kramer  
14  
15 146 HSD or Student's test. All statistical analyses were performed with JMP 8 software (SAS Institute  
16  
17 147 Inc., Cary, NC, USA).  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

148 **3. Results and Discussion**

149 **3.1 Effect of US processing on physicochemical and antioxidant parameters**

150 The colour of a food is an important freshness-related attribute for consumers and colour changes  
151 in a food product may affect their overall acceptability (Pingret et al. 2013). Previous studies  
152 suggested that US processing could affect the colour attributes of fruit and vegetables (Alexandre  
153 et al. 2012; Fava et al. 2011). However, in the current study, no significant differences were  
154 observed in colour parameters of *calçot* samples after sonication (Table 1). Birmpa, Sfika, &  
155 Vantarakis (2013) reported significant colour changes in lettuce leaves after US processing (37  
156 kHz, 30 W L<sup>-1</sup>) for 30, 45, or 60 min. The authors of that study suggested that a significant non-  
157 enzymatic browning could be responsible for the observed colour changes. The  $\Delta E^*$  combines  
158 the change in  $L^*$ ,  $a^*$ , and  $b^*$  values to quantify the colour deviation from a standard reference  
159 sample. Those samples with  $\Delta E^* > 3$  display a visible colour deviation (Wibowo et al. 2015). As  
160 expected, and shown in Table 1, US-treated *calçots* showed a  $\Delta E^* < 3$ . Moreover, BI values of  
161 all samples were similar and there were no significant differences ( $p < 0.05$ ) among them. Similar  
162 results were obtained previously after US processing (40 kHz, 500 W) of strawberries (do Rosário  
163 et al. 2017).

164 In addition, appearance and texture changes are two key characteristics determining the  
165 acceptability of fresh-cut fruit and vegetables (Toivonen and Brummell 2008). The texture of a  
166 food treated by US can be determined by the structure changes of proteins and enzymes during  
167 sonication (de São José et al. 2014). In the current study, as shown in Table 1, no significant  
168 differences were observed between the firmness and weight of the control and US-treated samples  
169 ( $p < 0.05$ ). Results were comparable to those previously reported by Ding et al. (2015), who  
170 observed that the firmness of strawberries after US (40 kHz, 240 W) treatment for 10 min did not  
171 change significantly. In addition, Alexandre et al. (2012) observed a higher firmness retention (16  
172 %) in US-treated (2 min,  $15 \pm 2$  °C, 35 kHz, 120 W) strawberries when compared to water-washed  
173 strawberries.

174 Besides physical attributes of foods such as colour or firmness, US treatment could affect minor  
175 components associated with TAC and phytochemical content. In the current study, two methods,  
176 DPPH and FRAP, were used to investigate the changes in total TAC of *calçots* after US treatment.  
177 Antioxidant capacity of *calçots* before and after processing are shown in Figure 1. Although  
178 higher treatment times resulted in a significant decrease in the TAC of the samples (data not  
179 shown), US processing for either 10, 25, or 45 min had no effect on the TAC of *calçots* ( $p<0.05$ ).  
180 Results obtained using the FRAP were in line with those obtained using the DPPH method.  
181 Results obtained herein were in agreement with those reported by Wang et al. (2015) who showed  
182 that US treatment (8 min, 25 °C, 20 kHz, 106.19 W L<sup>-1</sup>) had no effect on the TAC of cherry  
183 tomatoes. Similar results were also reported after processing of eggplant (Colucci et al. 2018).  
184 However, Muzaffar et al. (2016) and Gani et al. (2016) recently reported an increase of TAC in  
185 US-treated (25 °C, 33 kHz, 60 W) at different times (0, 10, 20, 30, 40 and 60 min) cherries and  
186 strawberries when compared to the untreated samples.

187 The TPC of the control and US-treated *calçots* is shown in Figure 1. In the current study, treating  
188 for either 10, 25, or 45 min did not affect the TPC of the samples when compared to the untreated  
189 control ( $p<0.05$ ). Results were in agreement with those obtained by Santos et al. (2015) who  
190 reported that both TAC and TPC of fresh-cut mango were maintained after US processing (25 °C,  
191 25 kHz, 55 W L<sup>-1</sup>) for 30 min. Previous authors observed a decrease in the TPC of US-treated  
192 fruit and vegetables caused by a oxidation due to hydroxyl radicals formed by cavitation (de São  
193 José et al. 2014; Rawson et al. 2011). However, Yeoh & Ali (2017) showed that the TPC of fresh-  
194 cut pineapple was increased after processing at 25 and 29 W for 10-15 min. The calculated TPC  
195 of the untreated and US-treated *calçots* correlates well with the observed TAC before and after  
196 processing.

### 197 **3.2 Effect of thermal processing on the nutritional quality of *calçots***

198 *Calçots* are generally eaten cooked after roasting. However, vitamins, phenolic compounds, and  
199 other health-promoting compounds have been shown to be heavily lost during thermal processing

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

200 (Kapusta-Duch et al. 2016; Soares et al. 2017). The effects of thermal processing on the TAC and  
201 TPC of *calçots* are shown in Figure 1. Overall, TAC of all samples increased after roasting  
202 ( $p<0.05$ ). In the same way, Juárez et al. (2016) reported that TAC of chopped onions increased  
203 after cooking (150 °C for 10 min + 110 °C for 5 min). In summary, the increase of TAC after  
204 roasting (270 °C, 8 min) could be due to: (1) liberation of high amount of antioxidant compounds  
205 due to thermal destruction of cell walls and sub cellular compartments; (2) production of  
206 antioxidant compounds with high radical scavenging activity; (3) suppression of oxidation  
207 capacity of antioxidant compounds due to the thermal inactivation of oxidative enzymes; (4)  
208 production of new no-nutrient antioxidants or the formation of new compounds such as Maillard  
209 reactions' compounds which could have antioxidant activity (Jiménez-Monreal et al. 2009;  
210 Morales and Babbel 2002).

211 Moreover, there were no significant differences ( $p>0.05$ ) between TPC of fresh and roasted  
212 *calçots* (270 °C, 8 min) at each processing time. However, Sharma et al. (2015) reported that  
213 heating at 80 °C, 100 °C, and 120 °C for 30 min increased and at 150 °C for 30 min decreased  
214 the total phenolic content for all studied onion varieties. Furthermore, Guillén et al. (2017) showed  
215 that cooking (90-100 °C) reduced the initial phenolic content in broccoli, green beans, artichokes  
216 and carrots. Notwithstanding, Rawson et al. (2013) reported that the decrease observed in total  
217 phenolic content was higher in boiled (30 min) than in roasted (160 °C, 15 min) fennel slices.

### 218 **3.3 Effect of US processing on the microbiological quality of *calçots***

219 There are indications that suggest that US can be used in the food industry, alone or associated  
220 with chemical sanitizers, to remove dirt and food residues as well as to inactivate microorganisms  
221 from the surfaces of fruit and vegetables (de São José et al. 2014). Microbial inactivation occurs  
222 because of cavitation. In the current study, processing for 10 min did not significantly reduce the  
223 total aerobic count in the US-treated *calçots* when compared to the untreated samples (Figure 2).  
224 However, US processing for 45 min significantly reduced the microbial load (around 1.0-log) of  
225 the samples ( $p<0.05$ ). In all cases, the microbial load was not higher than 6 log (cfu g<sup>-1</sup>). Bilek &

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

226 Turantaş (2013) recently suggested that US processing for 10 min, alone or in combination with  
227 other strategy, is generally enough to decontaminate fruit and vegetables. Indeed, Ding et al.  
228 (2015) reported that US treatment (40 kHz, 240 W) for 10 min removed 0.71 log cfu g<sup>-1</sup> for total  
229 aerobic bacteria on cherry tomatoes. In the same way, Cao et al. (2010) observed that numbers of  
230 aerobic microorganism of strawberries decreased from 2.15 ± 0.02 to 1.49 ± 0.01 log<sub>10</sub> cfu g<sup>-1</sup>  
231 after US treatment (20 °C, 40 kHz, 350 W) for 10 min.

### 232 **3.4 Resistance of TAC and TPC to a simulated gastrointestinal digestion**

233 In reference to evaluation the biological activity of *calçots* is much more relevant to know TAC  
234 and TPC potentially available for further intestinal absorption and/or protection than the  
235 quantification in the food matrix (Carbonell-Capella et al. 2014). Results obtained herein  
236 suggested that the TAC and TPC were statistically lower after gastrointestinal digestion when  
237 compared to the control ( $p < 0.05$ ; Figure 3). Similar results were reported by Ramírez-Moreno  
238 et al. (2018), where TAC and TPC of blackberry juice treated with US (20 kHz, 1500 W) at  
239 different times (0, 15 and 25 min) and amplitudes (60 and 80 %) decreased drastically after *in*  
240 *vitro* digestion. Recent studies have evaluated the effect of US treatment on the bioaccessibility  
241 of other compounds such as lycopene. For example, Anese et al. (2013, 2015) studied the effect  
242 of US treatment on the bioaccessibility of lycopene of tomato pulp. Despite the high decrease  
243 observed in TAC values, control (0 min) and US-treated samples (10 and 25 min) presented lower  
244 decrease (around 60 %) than roasted samples (70-90 %). The same tendency was observed in TPC  
245 values and *calçots* (raw or roasted) treated for 10 min presented the lowest values (around 70 %).  
246 The observed differences could be due to the sensitivity and instability to the pH changes and  
247 enzymatic activity during *in vitro* digestion of antioxidant compounds formed in the thermal  
248 processing. In the recent study carried out by de Lima et al. (2017), the effect of three different  
249 cooking methods (boiling, steaming and microwave) on the bioaccessibility of TAC and TPC of  
250 cassava. In that study a drastic decrease of TAC and TPC after *in vitro* digestion was observed  
251 and the bioaccessibility was similar in all studied samples. Recent studies have evaluated the  
252 effect of different cooking treatment on the bioaccessibility of other compounds. For example,



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

253 (Palmero et al. 2014) studied the effect of thermal treatment on the bioaccessibility of  $\beta$ -carotene  
254 of orange carrots and lycopene of red carrots and tomatoes. The vast majority of research on  
255 roasting and subsequent digestion has been carried out with cereals or coffee/cacao beans (Ribas-  
256 Agustí et al. 2017).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

257 **4. Conclusions**

258 The physical and microbiological quality and antioxidant capacity of fresh-cut *calçots* after  
259 ultrasound treatment was measured and those samples were also roasted (270 °C, 8 min) and  
260 digested. Minimally processed *calçots* pre-treated with ultrasounds (40 kHz, 250 W) for 10, 25  
261 or 45 min retained colour, firmness and weight after processing. Ultrasound pre-treatment had no  
262 effect on the antioxidant properties of fresh-cut *calçots*, but both the thermal process (270 °C, 8  
263 min) and the *in vitro* digestion produced a considerable reduction. Although microbial load of all  
264 samples was lower than 6 log (cfu g<sup>-1</sup>), only a decrease could be observed in those samples treated  
265 for 45 min. Therefore, pre-treatment with ultrasound showed potential to be used as a  
266 complementary treatment in the food industry. It is necessary to emphasize that this study was a  
267 first step to optimize the treatment conditions. Additional studies into the effect of ultrasound on  
268 the enzymatic activity in this type of fresh-cut vegetables should be undertaken in future works.

269 **Acknowledgments**

1  
2  
3 270 This work was supported by ACCIÓ (Generalitat of Catalonia, RD14-1-004), Sociedad Agrícola  
4  
5 271 i Secció de Crèdit de Valls S.C.C.L., Cooperativa of Cambrils, and PGI ‘Calçot de Valls’. This  
6  
7 272 work was also supported by the ‘Secretaria d’Universitats i Recerca del Departament d’Economia  
8  
9 273 i Coneixement’ (FI-2017-B2-00164, L. Zudaire) and CERCA Programme of Generalitat de  
10  
11 274 Catalunya. T. Lafarga is in receipt of a ‘Juan de la Cierva’ contract awarded by the Spanish  
12  
13 275 Ministry of Economy, Industry, and Competitiveness (FJCI-2016-29541). I. Aguiló-Aguayo  
14  
15 276 thanks to the National Programme for the Promotion of Talent and Its Employability of the  
16  
17 277 ‘Ministerio de Economía, Industria y Competitividad’ of the Spanish Government and to the  
18  
19 278 European Social Fund for the Postdoctoral Senior Grant ‘Ramon y Cajal’ (RYC-2016-19949).  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

279 **References**

- 1  
2  
3 280 Aguiló-Aguayo, I., Simó, J., Ivars, N., Villaró, S., Zudaire, L., Echeverria, G., et al. (2015).  
4  
5 281 Suitability of the ‘calçots’ (*Allium cepa* L.) for minimal processing. In *2nd Euro-*  
6  
7 282 *Mediterranean Symposium on Fruit and Vegetable Processing*. Avignon, France.  
8  
9  
10 283 Alegre, I., Viñas, I., Usall, J., Anguera, M., & Abadias, M. (2011). Microbiological and  
11  
12 284 physicochemical quality of fresh-cut apple enriched with the probiotic strain *Lactobacillus*  
13  
14 285 *rhamnosus* GG. *Food microbiology*, 28(1), 59–66. doi:10.1016/j.fm.2010.08.006  
15  
16  
17  
18 286 Alexandre, E. M. C., Brandão, T. R. S., & Silva, C. L. M. (2012). Efficacy of non-thermal  
19  
20 287 technologies and sanitizer solutions on microbial load reduction and quality retention of  
21  
22 288 strawberries. *Journal of Food Engineering*, 108(3), 417–426.  
23  
24 289 doi:10.1016/j.jfoodeng.2011.09.002  
25  
26  
27  
28 290 Altisent, R., Plaza, L., Alegre, I., Viñas, I., & Abadias, M. (2014). Comparative study of improved  
29  
30 291 vs. traditional apple cultivars and their aptitude to be minimally processed as ‘ready to eat’  
31  
32 292 apple wedges. *LWT - Food Science and Technology*, 58(2), 541–549.  
33  
34 293 doi:10.1016/j.lwt.2014.03.019  
35  
36  
37  
38 294 Anese, M., Bot, F., Panozzo, A., Mirolo, G., & Lippe, G. (2015). Effect of ultrasound treatment,  
39  
40 295 oil addition and storage time on lycopene stability and *in vitro* bioaccessibility of tomato  
41  
42 296 pulp. *Food Chemistry*, 172, 685–691. doi:10.1016/j.foodchem.2014.09.140  
43  
44  
45 297 Anese, M., Mirolo, G., Beraldo, P., & Lippe, G. (2013). Effect of ultrasound treatments of tomato  
46  
47 298 pulp on microstructure and lycopene *in vitro* bioaccessibility. *Food Chemistry*, 136(2), 458–  
48  
49 299 463. doi:10.1016/j.foodchem.2012.08.013  
50  
51  
52  
53 300 Awad, T. S., Moharram, H. A., Shaltout, O. E., Asker, D., & Youssef, M. M. (2012). Applications  
54  
55 301 of ultrasound in analysis, processing and quality control of food: A review. *Food Research*  
56  
57 302 *International*. Elsevier B.V. doi:10.1016/j.foodres.2012.05.004  
58  
59  
60 303 Bilek, S. E., & Turantaş, F. (2013). Decontamination efficiency of high power ultrasound in the

1  
2  
3  
4  
5 304 fruit and vegetable industry, a review. *International Journal of Food Microbiology*.  
6  
7 305 doi:10.1016/j.ijfoodmicro.2013.06.028  
8  
9  
10 306 Birmpa, A., Sfika, V., & Vantarakis, A. (2013). Ultraviolet light and Ultrasound as non-thermal  
11  
12 307 treatments for the inactivation of microorganisms in fresh ready-to-eat foods. *International*  
13  
14 308 *Journal of Food Microbiology*, 167(1), 96–102. doi:10.1016/j.ijfoodmicro.2013.06.005  
15  
16  
17 309 Bouayed, J., Deußer, H., Hoffmann, L., & Bohn, T. (2012). Bioaccessible and dialysable  
18  
19 310 polyphenols in selected apple varieties following *in vitro* digestion vs. their native patterns.  
20  
21 311 *Food Chemistry*, 131(4), 1466–1472. doi:10.1016/j.foodchem.2011.10.030  
22  
23  
24 312 Cao, S., Hu, Z., Pang, B., Wang, H., Xie, H., & Wu, F. (2010). Effect of ultrasound treatment on  
25  
26 313 fruit decay and quality maintenance in strawberry after harvest. *Food Control*, 21(4), 529–  
27  
28 314 532. doi:10.1016/j.foodcont.2009.08.002  
29  
30  
31 315 Carbonell-Capella, J. M., Buniowska, M., Barba, F. J., Esteve, M. J., & Frígola, A. (2014).  
32  
33 316 Analytical methods for determining bioavailability and bioaccessibility of bioactive  
34  
35 317 compounds from fruits and vegetables: A review. *Comprehensive Reviews in Food Science*  
36  
37 318 *and Food Safety*, 13(2), 155–171. doi:10.1111/1541-4337.12049  
38  
39  
40 319 Chemat, F., Zill-E-Huma, & Khan, M. K. (2011). Applications of ultrasound in food technology:  
41  
42 320 Processing, preservation and extraction. *Ultrasonics Sonochemistry*, 18(4), 813–835.  
43  
44 321 doi:10.1016/j.ultsonch.2010.11.023  
45  
46  
47 322 Colás-Medà, P., Abadias, M., Altisent, R., Alegre, I., Plaza, L., Gilabert, V., et al. (2016).  
48  
49 323 Development of a fresh-cut product based on pears and the subsequent evaluation of Its shelf  
50  
51 324 life under commercial conditions and after a cold chain break. *Journal of Food and Nutrition*  
52  
53 325 *Research*, 4(9), 582–591. doi:10.12691/jfnr-4-9-4  
54  
55  
56 326 Colucci, D., Fissore, D., Rossello, C., & Carcel, J. A. (2018). On the effect of ultrasound-assisted  
57  
58 327 atmospheric freeze-drying on the antioxidant properties of eggplant. *Food Research*  
59  
60 328 *International*. doi:10.1016/j.foodres.2018.01.022  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

329 D.A.R.P. (2009). ORDRE AAR/414/2009, de 21 de setembre, per la qual s'aprova el Reglament  
330 de la Indicació Geogràfica Protegida Calçot de Valls.  
331 [http://portaljuridic.gencat.cat/ca/pjur\\_ocults/pjur\\_resultats\\_fitxa/?documentId=503886&la](http://portaljuridic.gencat.cat/ca/pjur_ocults/pjur_resultats_fitxa/?documentId=503886&language=ca_ES&action=fitxa)  
332 [nguage=ca\\_ES&action=fitxa](http://portaljuridic.gencat.cat/ca/pjur_ocults/pjur_resultats_fitxa/?documentId=503886&language=ca_ES&action=fitxa)

333 de Lima, A. C. S., da Rocha Viana, J. D., de Sousa Sabino, L. B., da Silva, L. M. R., da Silva, N.  
334 K. V., & de Sousa, P. H. M. (2017). Processing of three different cooking methods of  
335 cassava: Effects on *in vitro* bioaccessibility of phenolic compounds and antioxidant activity.  
336 *LWT - Food Science and Technology*, 76, 253–258. doi:10.1016/j.lwt.2016.07.023

337 de São José, J., de Andrade, N. J., Ramos, A. M., Vanetti, M., Stringheta, P., & Chaves, J. (2014).  
338 Decontamination by ultrasound application in fresh fruits and vegetables. *Food Control*, 45,  
339 36–50. doi:10.1016/j.foodcont.2014.04.015

340 Ding, T., Ge, Z., Shi, J., Xu, Y. T., Jones, C. L., & Liu, D. H. (2015). Impact of slightly acidic  
341 electrolyzed water (SAEW) and ultrasound on microbial loads and quality of fresh fruits.  
342 *LWT - Food Science and Technology*, 60(2), 1195–1199. doi:10.1016/j.lwt.2014.09.012

343 do Rosário, D. K. A., da Silva Mutz, Y., Peixoto, J. M. C., Oliveira, S. B. S., de Carvalho, R. V.,  
344 Carneiro, J. C. S., et al. (2017). Ultrasound improves chemical reduction of natural  
345 contaminant microbiota and *Salmonella enterica* subsp. *enterica* on strawberries.  
346 *International Journal of Food Microbiology*, 241, 23–29.  
347 doi:10.1016/j.ijfoodmicro.2016.10.009

348 EC No 905/2002. Commission Regulation (EC) No 905/2002 of 30 May 2002 supplementing the  
349 Annex to Regulation (EC) No 2400/96 on the entry of certain names in the 'Register of  
350 protected designations of origin and protected geographical indications' [2002] OJ L 142/27.

351 Fava, J., Hodara, K., Nieto, A., Guerrero, S., Alzamora, S. M., & Castro, M. A. (2011). Structure  
352 (micro, ultra, nano), color and mechanical properties of *Vitis labrusca* L. (grape berry) fruits  
353 treated by hydrogen peroxide, UV-C irradiation and ultrasound. *Food Research*

- 354 *International*, 44(9), 2938–2948. doi:10.1016/j.foodres.2011.06.053
- 1  
2  
3 355 Gani, A., Baba, W. N., Ahmad, M., Shah, U., Khan, A. A., Wani, I. A., et al. (2016). Effect of  
4  
5 356 ultrasound treatment on physico-chemical, nutraceutical and microbial quality of  
6  
7 357 strawberry. *LWT - Food Science and Technology*, 66, 496–502.  
8  
9 358 doi:10.1016/j.lwt.2015.10.067
- 11  
12 359 Guillén, S., Mir-Bel, J., Oria, R., & Salvador, M. L. (2017). Influence of cooking conditions on  
13  
14 360 organoleptic and health-related properties of artichokes, green beans, broccoli and carrots.  
15  
16 361 *Food Chemistry*, 217, 209–216. doi:10.1016/j.foodchem.2016.08.067
- 18  
19  
20 362 Jiménez-Monreal, A. M., García-Diz, L., Martínez-Tomé, M., Mariscal, M., & Murcia, M. A.  
21  
22 363 (2009). Influence of cooking methods on antioxidant activity of vegetables. *Journal of Food*  
23  
24 364 *Science*, 74(3), 97–103. doi:10.1111/j.1750-3841.2009.01091.x
- 26  
27  
28 365 Juárez, I., Ludwig, I. A., Huarte, E., Pereira-Caro, G., Moreno-Rojas, J. M., Cid, C., & De Peña,  
29  
30 366 M. P. (2016). Influence of heat treatment on antioxidant capacity and (poly)phenolic  
31  
32 367 compounds of selected vegetables. *Food Chemistry*, 197, 466–473.  
33  
34 368 doi:10.1016/j.foodchem.2015.10.139
- 36  
37  
38 369 Kapusta-Duch, J., Kusznierevicz, B., Leszczyńska, T., & Borczak, B. (2016). Effect of cooking  
39  
40 370 on the contents of glucosinolates and their degradation products in selected *Brassica*  
41  
42 371 vegetables. *Journal of Functional Foods*, 23, 412–422. doi:10.1016/j.jff.2016.03.006
- 44  
45 372 Kentish, S., & Ashokkumar, M. (2011). The physical and chemical effects of ultrasound. In H.  
46  
47 373 Feng, G. V. Barbosa-Cánovas, & J. Weiss (Eds.), *Ultrasound technologies for food and*  
48  
49 374 *bioprocessing* (pp. 1–12). New York: Springer.
- 51  
52  
53 375 Lagnika, C., Zhang, M., & Mothibe, K. J. (2013). Effects of ultrasound and high pressure argon  
54  
55 376 on physico-chemical properties of white mushrooms (*Agaricus bisporus*) during postharvest  
56  
57 377 storage. *Postharvest Biology and Technology*, 82, 87–94.  
58  
59 378 doi:10.1016/j.postharvbio.2013.03.006
- 60  
61  
62  
63  
64  
65

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- 379 Liu, C., Ma, T., Hu, W., Tian, M., & Sun, L. (2016). Effects of aqueous ozone treatments on  
380 microbial load reduction and shelf life extension of fresh-cut apple. *International Journal*  
381 *of Food Science and Technology*, 51(5), 1099–1109. doi:10.1111/ijfs.13078
- 382 Minekus, M., Alminger, M., Alvito, P., Ballance, S., Bohn, T., Bourlieu, C., et al. (2014). A  
383 standardised static *in vitro* digestion method suitable for food - an international consensus.  
384 *Food & function*, 5(6), 1113–24. doi:10.1039/c3fo60702j
- 385 Morales, F. J., & Babbel, M.-B. (2002). Antiradical efficiency of Maillard reaction mixtures in a  
386 hydrophilic media. *Journal of agricultural and food chemistry*, 50(10), 2788–2792.  
387 doi:10.1021/jf011449u
- 388 Muzaffar, S., Ahmad, M., Wani, S. M., Gani, A., Baba, W. N., Shah, U., et al. (2016). Ultrasound  
389 treatment: effect on physicochemical, microbial and antioxidant properties of cherry  
390 (*Prunus avium*). *Journal of Food Science and Technology*, 53(6), 2752–2759.  
391 doi:10.1007/s13197-016-2247-3
- 392 Palmero, P., Lemmens, L., Hendrickx, M., & Van Loey, A. (2014). Role of carotenoid type on  
393 the effect of thermal processing on bioaccessibility. *Food Chemistry*, 157, 275–282.  
394 doi:10.1016/j.foodchem.2014.02.055
- 395 Pingret, D., Fabiano-Tixier, A. S., & Chemat, F. (2013). Degradation during application of  
396 ultrasound in food processing: A review. *Food Control*, 31(2), 593–606.  
397 doi:10.1016/j.foodcont.2012.11.039
- 398 Piyasena, P., Mohareb, E., & McKellar, R. C. (2003). Inactivation of microbes using ultrasound:  
399 A review. *International Journal of Food Microbiology*, 87(3), 207–216.  
400 doi:10.1016/S0168-1605(03)00075-8
- 401 Plaza, L., Altisent, R., Alegre, I., Viñas, I., & Abadias, M. (2016). Changes in the quality and  
402 antioxidant properties of fresh-cut melon treated with the biopreservative culture  
403 *Pseudomonas graminis* CPA-7 during refrigerated storage. *Postharvest Biology and*



404 *Technology*, 111, 25–30. doi:10.1016/j.postharvbio.2015.07.023

405 Ramírez-Moreno, E., Zafra-Rojas, Q. Y., Arias-Rico, J., Ariza-Ortega, J. A., Alanís-García, E.,  
406 & Cruz-Cansino, N. (2018). Effect of ultrasound on microbiological load and antioxidant  
407 properties of blackberry juice. *Journal of Food Processing and Preservation*, 42(2), 1–6.  
408 doi:10.1111/jfpp.13489

409 Rawson, A., Hossain, M. B., Patras, A., Tuohy, M., & Brunton, N. (2013). Effect of boiling and  
410 roasting on the polyacetylene and polyphenol content of fennel (*Foeniculum vulgare*) bulb.  
411 *Food Research International*, 50(2), 513–518. doi:10.1016/j.foodres.2011.01.009

412 Rawson, A., Tiwari, B. K., Patras, A., Brunton, N., Brennan, C., Cullen, P. J., & O'Donnell, C.  
413 (2011). Effect of thermosonication on bioactive compounds in watermelon juice. *Food*  
414 *Research International*, 44(5), 1168–1173. doi:10.1016/j.foodres.2010.07.005

415 Ribas-Agustí, A., Martín-Belloso, O., Soliva-Fortuny, R., & Elez-Martínez, P. (2017). Food  
416 processing strategies to enhance phenolic compounds bioaccessibility and bioavailability in  
417 plant-based foods. *Critical Reviews in Food Science and Nutrition*, 1–18.  
418 doi:10.1080/10408398.2017.1331200

419 Santos, J. G., Fernandes, F. A. N., de Siqueira Oliveira, L., & de Miranda, M. R. A. (2015).  
420 Influence of ultrasound on fresh-cut mango quality through evaluation of enzymatic and  
421 oxidative metabolism. *Food and Bioprocess Technology*, 8(7), 1532–1542.  
422 doi:10.1007/s11947-015-1518-8

423 Sharma, K., Ko, E. Y., Assefa, A. D., Ha, S., Nile, S. H., Lee, E. T., & Park, S. W. (2015).  
424 Temperature-dependent studies on the total phenolics, flavonoids, antioxidant activities, and  
425 sugar content in six onion varieties. *Journal of Food and Drug Analysis*, 23(2), 243–252.  
426 doi:10.1016/j.jfda.2014.10.005

427 Simó, J., Valero, J., Plans, M., Romero del Castillo, R., & Casañas, F. (2013). Breeding onions  
428 (*Allium cepa* L.) for consumption as ‘calçots’ (second-year resprouts). *Scientia*

429 *Horticulturae*, 152, 74–79. doi:10.1016/j.scienta.2013.01.011

430 Singleton, V., Orthofer, R., & Lamuela-Raventós, R. M. (1999). Analysis of total phenols and  
431 other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent.  
432 *METHODS IN ENZIMOLOGY*, 299, 152–178.

433 Soares, A., Carrascosa, C., & Raposo, A. (2017). Influence of different cooking methods on the  
434 concentration of glucosinolates and vitamin C in broccoli. *Food and Bioprocess  
435 Technology*, 10(8), 1387–1411. doi:10.1007/s11947-017-1930-3

436 Toivonen, P. M. A., & Brummell, D. A. (2008). Biochemical bases of appearance and texture  
437 changes in fresh-cut fruit and vegetables. *Postharvest Biology and Technology*, 48(1), 1–14.  
438 doi:10.1016/j.postharvbio.2007.09.004

439 Wang, W., Ma, X., Zou, M., Jiang, P., Hu, W., Li, J., et al. (2015). Effects of ultrasound on  
440 spoilage microorganisms, quality, and antioxidant capacity of postharvest cherry tomatoes.  
441 *Journal of Food Science*, 80(10), C2117–C2126. doi:10.1111/1750-3841.12955

442 Welti-Chanes, J., Morales-de la Peña, M., Jacobo-Velázquez, D. A., & Martín-Belloso, O. (2017).  
443 *Opportunities and challenges of ultrasound for food processing: An industry point of view.*  
444 *Ultrasound: Advances for food processing and preservation.* Academic Press.  
445 doi:10.1016/B978-0-12-804581-7.00019-1

446 Wibowo, S., Vervoort, L., Tomic, J., Santiago, J. S., Lemmens, L., Panozzo, A., et al. (2015).  
447 Colour and carotenoid changes of pasteurised orange juice during storage. *Food Chemistry*,  
448 171, 330–340. doi:10.1016/j.foodchem.2014.09.007

449 Yeoh, W. K., & Ali, A. (2017). Ultrasound treatment on phenolic metabolism and antioxidant  
450 capacity of fresh-cut pineapple during cold storage. *Food Chemistry*, 216, 247–253.  
451 doi:10.1016/j.foodchem.2016.07.074

452 Zudaire, L., Viñas, I., Abadias, M., Simó, J., Echeverria, G., Plaza, L., & Aguiló-Aguayo, I.  
453 (2017). Quality and bioaccessibility of total phenols and antioxidant activity of *calçots*

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

454 (*Allium cepa* L.) stored under controlled atmosphere conditions. *Postharvest Biology and*  
455 *Technology*, 129, 118–128. doi:10.1016/j.postharvbio.2017.03.013

456

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

457 **Figure Captions**

458 **Fig. 1. Effect of US and thermal processing on the TAC measured using the DPPH (A) and**  
459 **FRAP (B) methods and on the TPC (C) of US- and thermally-treated calçots.**


460 Lower case letters indicate significant differences between fresh samples (black bars) and capital  
461 letters indicate significant differences between roasted (270 °C, 8 min) samples (grey bars). \*  
462 indicates significant differences between fresh and roasted samples. The criterion for statistical  
463 significance was  $p<0.05$ . The error bars represent the standard errors of the mean of three  
464 independent measurements.


465 **Fig. 2. Effect of US processing on the total aerobic count of fresh-cut calçots.**


466 Lower case letters indicate significant differences between samples. The criterion for statistical  
467 significance was  $p<0.05$ . The error bars represent standard errors of the mean of independent  
468 measurements.

469 **Fig. 3. Resistance of TAC assessed using the DPPH (A) and FRAP (B) method and TPC (C)**  
470 **of US- and thermally-treated calçots to a simulated gastrointestinal digestion.**

471 Lower case letters indicate significant differences between samples (grey bars) after *in vitro*  
472 simulated digestion. \* indicates significant differences between undigested (black bars) and  
473 digested samples (grey bars). The criterion for statistical significance was  $p<0.05$ . The error bars  
474 represent the standard errors of the mean of three independent measurements







**Table 1.** Colour parameters, firmness, and weight of untreated and US-treated *calçots* (fresh). Values represent the means of independent experiments  $\pm$  standard deviation. Different letters in the same column indicate significant differences between samples ( $p < 0.05$ ).

Sample	$h^\circ$	BI	$\Delta E^*$	Firmness (N)	Weight (g)
0 min (control)	$104.54 \pm 3.31^a$	$7.50 \pm 1.87^a$	-	$138.00 \pm 36.94^a$	$52.89 \pm 14.04^a$
10 min	$103.46 \pm 2.28^a$	$7.41 \pm 2.20^a$	$4.70 \pm 3.08^a$	$123.08 \pm 41.80^a$	$54.73 \pm 13.91^a$
25 min	$105.79 \pm 3.19^a$	$6.41 \pm 1.65^a$	$5.93 \pm 4.86^a$	$102.08 \pm 33.73^a$	$51.23 \pm 16.56^a$
45 min	$105.00 \pm 2.78^a$	$7.28 \pm 1.77^a$	$4.04 \pm 2.48^a$	$121.95 \pm 30.93^a$	$56.27 \pm 15.17^a$



1    **Abstract**

2    The effect of ultrasound (US) treatment (40 kHz, 250 W) for 0, 10, 25 and 45 min on the physical  
3    and microbiological quality, total antioxidant capacity (TAC) and total phenolic content (TPC) of  
4    *calçots* (*Allium cepa* L.) was evaluated. Moreover, the effect of roasting (270 °C, 8 min) and *in*  
5    *vitro* simulated digestion on the antioxidant properties was studied. Overall, US treatment had no  
6    effect of the physical quality and antioxidant properties of *calçots* regardless the treatment time,  
7    while thermal processing produced an increase on the TAC and maintenance in TPC.  
8    Furthermore, the digestion process caused a remarkable decrease on the TAC and TPC, but that  
9    decrease was higher in roasted than in fresh samples. The microbial load of all US-treated fresh  
10   samples was below 6 log (cfu g<sup>-1</sup>) and a decrease of 1-log reduction was observed after treating  
11   for 45 min. Those results indicated that US pre-treatment had no negative effects on the quality  
12   of *calçot* while produced a decrease on the microbial load at high processing times.

13   **Keywords:** *Allium cepa* L.; thermal processing; gastrointestinal digestion; antioxidant activity;  
14   novel technologies