



A characterization of the generalized Liénard polynomial differential systems having invariant algebraic curves

Jaume Giné^{a,*}, Jaume Llibre^b

^a Departament de Matemàtica, Universitat de Lleida, Avda. Jaume II, 69, 25001 Lleida, Catalonia, Spain

^b Departament de Matemàtiques, Universitat Autònoma de Barcelona, 08193 Bellaterra, Barcelona, Catalonia, Spain

ARTICLE INFO

Article history:

Received 13 December 2021
Received in revised form 10 March 2022
Accepted 31 March 2022
Available online 25 April 2022

Keywords:

Liénard polynomial differential systems
Invariant algebraic curve
First integrals

ABSTRACT

The generalized Liénard polynomial differential systems are the differential systems of the form $x' = y$, $y' = -f(x)y - g(x)$, where f and g are polynomials.

We characterize all the generalized Liénard polynomial differential systems having an invariant algebraic curve. We show that the first four higher coefficients of the polynomial in the variable y , defining the invariant algebraic curve, determine completely the generalized Liénard polynomial differential system. This fact does not hold for arbitrary polynomial differential systems.

2010 mathematics subject classification: Primary 34A05. Secondary 34C05, 37C10.

© 2022 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction and statement of the main results

In this work we study the generalized Liénard polynomial differential systems of the form

$$x' = y, \quad y' = -f(x)y - g(x), \quad (1)$$

where the degrees of the polynomials f and g are m and n respectively.

Let $F(x, y)$ be a polynomial such that

$$\frac{\partial F}{\partial x}y + \frac{\partial F}{\partial y}(-f(x)y - g(x)) = KF, \quad (2)$$

for some polynomial $K = K(x, y)$. Then $F(x, y) = 0$ is an *invariant algebraic curve* of the differential system (1), i.e. if an orbit of system (1) has a point on the curve $F(x, y) = 0$, the whole orbit is contained in this curve. The polynomial K is called as/or to be the *cofactor* of the invariant algebraic curve $F(x, y) = 0$.

The knowledge of the algebraic curves of system (1) allows to study the Darboux and Liouvillian theories of integrability, see [1–4] and references therein. In fact the existence of invariant algebraic curves is a measure of the integrability in such theories. Another problem is finding a bound for the degree of the irreducible invariant algebraic curves of system (1). This problem goes back to Poincaré for any polynomial differential system and it is known as the *Poincaré problem* for the invariant algebraic curves. The invariant algebraic curves of generalized Liénard

systems (1) have been studied by several authors in function of degrees of f and g , see for instance [5–12] and references therein. The determination of invariant algebraic curves is also important when we study the algebraic limit cycles of such systems, see [13–15]. Several works are also devoted to the Liouville integrability of such systems, see [16–20]. Finally we remark that a new method to determine the invariant algebraic curves have been developed on [21–24] based on the solutions of the differential system expressed in Puiseux series. In this note we study the reciprocal problem. This problem consists in given an invariant algebraic curve characterize the generalized Liénard polynomial differential systems having such an invariant algebraic curve.

In the following we use the notation $a_j^k(x)$ to denote $(a_j(x))^k$.

Theorem 1. Assume that a generalized Liénard polynomial differential system (1) with the polynomials f and g non-identically zero has an invariant algebraic curve that we write as

$$F(x, y) = \sum_{j=0}^s a_j(x)y^{s-j} = 0 \text{ with } a_0(x) \neq 0 \text{ and } s \geq 2. \quad (3)$$

Then the polynomials f and g are

$$f(x) = \frac{sa_3(x) - (s-1)a_1(x)a_2(x) + (s-1)a_1^2(x)a_1'(x) - sa_2(x)a_1'(x)}{(s-1)a_1^2(x) - 2sa_2(x)},$$

$$g(x) = \frac{a_1(x)a_2(x)a_1'(x) + a_1(x)a_2(x) - 2a_2(x)a_2'(x)}{(s-1)a_1^2(x) - 2sa_2(x)},$$

$a_0(x)$ is a constant and the cofactor of $F(x, y) = 0$ only depends on x .

Theorem 1 is proved in Section 2.

* Corresponding author.

E-mail addresses: jaume.gine@udl.cat (J. Giné), jllibre@mat.uab.cat (J. Llibre).

Note that the common denominator in the expressions of $f(x)$ and $g(x)$ given in the statement of **Theorem 1** must divide their numerators, otherwise $f(x)$ and $g(x)$ would not be polynomials.

We remark that if a generalized Liénard polynomial differential system (1) has an invariant algebraic curve (3) the coefficients $a_1(x)$, $a_2(x)$ and $a_3(x)$ determine completely such differential system. Of course this is not true for general polynomial differential systems, for instance the polynomial differential system

$$x = a(x, y)F - c(x, y)\frac{\partial F}{\partial y}, \quad y = b(x, y)F + c(x, y)\frac{\partial F}{\partial x},$$

where a , b and c are arbitrary polynomials, has $F = F(x, y) = 0$ as an invariant algebraic curve.

Corollary 2. Under the assumptions of **Theorem 1** if the common denominator of the expressions of $f(x)$ and $g(x)$ are zero, then the polynomials $f(x)$ and $g(x)$ become

$$f(x) = \frac{a_1'(x)}{2}, \quad g(x) = \frac{a_2'(x)}{s} - \frac{a_1(x)a_1'(x)}{2s}.$$

Proposition 3. Under the assumptions of **Theorem 1** if $a_i(x) = \sum_{j=1}^3 a_{ij}x^j$ for $j=1, 2, 3$ are arbitrary polynomials such that the maximum degree of all them is 3, then the generalized Liénard polynomial differential systems having an irreducible invariant algebraic curve of degree 3 in the variable y are the following ones:

- I $f(x) = \frac{4a_{11}}{9}, \quad g(x) = -\frac{a_{11}}{27}(a_{10} + a_{11}x),$
 $F(x, y) = \frac{1}{27} \left((a_{10} + a_{11}x)(3a_{20} + a_{11}x(2a_{10} + a_{11}x)) \right.$
 $\quad + 9(3a_{20} + a_{11}x(2a_{10} + a_{11}x))y$
 $\quad \left. + 27(a_{10} + a_{11}x)y^2 + 27y^3 \right).$
- II $f(x) = \frac{5a_{11}}{9}, \quad g(x) = -\frac{2a_{11}}{27}(3a_{10} + a_{11}x),$
 $F(x, y) = \frac{1}{81} (3a_{10} + a_{11}x) (a_{10} + 3a_{11}x)$
 $\quad + \frac{1}{9} (3a_{10} + a_{11}x)(a_{10} + 3a_{11}x)y + (a_{10} + a_{11}x)y^2 + y^3.$
- III $f(x) = \frac{a_{11}}{3}, \quad g(x) = -\frac{2a_{11}^2}{9}x,$
 $F(x, y) = \frac{1}{27} (27a_{30} - 4a_{11}^3x^3 + 27a_{11}xy^2 + 27y^3).$
- IV $f(x) = \frac{2a_{11}^3 - k - 6a_{11}a_{22}}{3(a_{11}^2 - 3a_{22})}, \quad g(x) = \frac{a_{11}^4 + ka_{11} - 9a_{11}^2a_{22} + 18a_{22}^2}{9(a_{11}^2 - 3a_{22})},$
 $F(x, y) = \frac{1}{27} (-2a_{11}^3x^3 - 2kx^3 + 9a_{11}a_{22}x^3 + 27a_{22}x^2y + 27a_{11}xy^2 + 27y^3),$ where
 $k = (a_{11}^2 - 3a_{22})^{3/2}.$

Taking into account that the degrees of the polynomials f and g are m and n respectively, from **Corollary 2** it follows that in this case the $\deg a_1 = m + 1$ and $\deg a_2 \leq \max \{n, m(m + 1)\} + 1$.

Under the assumptions of **Corollary 2** we note that the generalized Liénard polynomial differential system (1) having an invariant algebraic curve (3) already the coefficients $a_1(x)$ and $a_2(x)$ determine completely such differential system.

Proposition 4. Under the assumptions of **Corollary 2** if $a_1(x) = \sum_{i=1}^4 a_{1i}x^i$ and $a_2(x) = \sum_{i=1}^4 a_{2i}x^i$ are arbitrary polynomials such that the maximum degree of both is 4, then the generalized Liénard polynomial differential

systems having an irreducible invariant algebraic curve of degree 4 in the variable y are the following ones:

- I $f(x) = \frac{a_{11}}{2}, \quad g(x) = -\frac{a_{11}^2}{8}x,$
 $F(x, y) = -\frac{a_{11}^4}{128}x^4 + \frac{a_{11}^3}{16}x^3y + a_{11}xy^3 + y^4$
- II $f(x) = \frac{a_{11}}{2}, \quad g(x) = -\frac{a_{11}}{8}(a_{10} + a_{11}x),$
 $F(x, y) = -\frac{a_{10}^4}{128} - \frac{a_{11}}{128}x(4a_{10}^3 + 6a_{10}^2a_{11}x + 4a_{10}a_{11}^2x^2 + a_{11}^3x^3)$
 $\quad + \frac{1}{16}(a_{10} + a_{11}x)^3y + (a_{10} + a_{11}x)y^3 + y^4$
- III $f(x) = \frac{1}{2}(a_{11} + 2a_{12}x + 3a_{13}x^2 + 4a_{14}x^3)$
 $g(x) = -\frac{1}{4}f(x)(a_{10} + a_{11}x + a_{12}x^2 + a_{13}x^3 + a_{14}x^4),$
 $F(x, y) = -\frac{1}{128} \left\{ a_{10} + x(a_{11} + x(a_{12} + x(a_{13} + a_{14}x))) \right\}^4 + \frac{1}{16}a_{10}$
 $\quad + x(a_{11} + x(a_{12} + x(a_{13} + a_{14}x)))^3y$
 $\quad + \left\{ a_{10} + x(a_{11} + x(a_{12} + x(a_{13} + a_{14}x))) \right\} y^3 + y^4.$

Corollary 2 and **Propositions 3** and **4** are proved in **Section 2**.

We remark that the next polynomial Liénard differential system shows that there are invariant algebraic curves of arbitrary degree in the variable y . In [25] the authors proved that the linear polynomial Liénard differential system

$$x = y, \quad y = x - \frac{p-1}{\sqrt{p}}y,$$

with $p > 1$ has the first integral $H = (\sqrt{p}y - x)(\sqrt{p}y + x)^p$. Therefore such differential system has invariant algebraic curves of degree $p + 1$ for all p .

2. Proofs

Proof of Theorem 1. If $F(x, y) = 0$ is an invariant algebraic curve of system (1) it must satisfy (2) with a cofactor of the form $K = \sum_{j=0}^s K_j(x)y^j$. From Eq. (2) it follows easily that $r \leq 1$. If $r = 1$ then the coefficient of y^{s+1} is $a_0'(x) - a_0(x)K_1(x) = 0$. Since $a_0(x)$ must be a polynomial it follows that $K_1(x) = 0$ and $a_0(x)$ is a constant. Then without loss of generality we can take $a_0(x) = 1$, because we can divide the invariant algebraic curve by the non-zero constant $a_0(x)$. In summary the cofactor of $F(x, y) = 0$ only depends on the variable x .

From the coefficient of y^s in Eq. (2) we obtain that the cofactor $K_0(x) = a_1'(x) - sf(x)$. And from the coefficient of y^{s-1} in Eq. (2) we get that

$$g(x) = \frac{1}{s} (a_2'(x) - (s-1)a_1(x)f(x) - K_0(x)a_1(x)).$$

Substituting $K_0(x)$ is the above expression we have

$$g(x) = \frac{1}{s} (a_2'(x) + a_1(x)f(x) - a_1(x)a_1'(x)). \tag{4}$$

The coefficient of y^{s-2} in Eq. (2) is

$$a_3'(x) - (s-2)a_2(x)f(x) - \frac{s-1}{s}a_1(x)(a_2'(x) + a_1(x)f(x) - a_1(x)a_1'(x))$$

$$= a_1'(x)a_2(x) - sa_2(x)f(x).$$

Consequently we obtain the function $f(x)$ stated in [Theorem 1](#), and substituting $f(x)$ in (4) we get the expression of $g(x)$ stated in the theorem. \square

Example 1. Consider the polynomial Liénard differential system

$$x = y, \quad y = x - \frac{1}{16}a^2(x^2 - 4)x^3 - a(x^2 - 1)y, \tag{5}$$

having the invariant algebraic curve

$$F(x, y) = y^2 + a\left(\frac{1}{2}x^3 - 2x\right)y - 4 + (a^2 + 1)x^2 - \frac{a^2x^4}{2} + \frac{a^2x^6}{16} \\ = y^2 + a_1(x)y + a_2(x).$$

Substituting $a_1(x)$ and $a_2(x)$ in the expressions of the polynomials f and g given in the statement of [Theorem 1](#) we obtain

$$f(x) = a(x^2 - 1), \quad g(x) = -x + \frac{1}{16}a^2(x^2 - 4)x^3.$$

Proof of Corollary 2. Since the denominator of the polynomial $f(x)$ divides its numerator, then $f(x)$ must vanish when $a_2(x) = (s - 1)a_1^2(x)/(2s)$. This implies that

$$s^2a_1^2(x)a_1'(x) - 3sa_1^2(x)a_1'(x) + 2a_1^2(x)a_1'(x) - 2s^2a_3'(x) = 0.$$

From this equation we get

$$a_3(x) = \frac{(s-2)(s-1)a_1^3(x)}{6s^2} + c_1,$$

where c_1 is an arbitrary constant. Substituting $a_3(x)$ in $f(x)$ we obtain

$$f(x) = \frac{N(x)}{D(x)} \tag{6}$$

where $N(x) = -2s^2a_2(x)a_1'(x) + 3s^2a_1^2(x)a_1'(x) - 5sa_1^2(x)a_1'(x) + 2a_1^2(x)a_1'(x) - 2s^2a_1(x)a_2'(x) + 2sa_1(x)a_2'(x)$ and $D(x) = 2s[(s - 1)a_1^2(x) - 2sa_2(x)]$.

Dividing the numerator $N(x)$ by the denominator $D(x)$ the quotient is $a_1(x)/2$, and the remainder $R(x)$ is equal to $2(s-1)a_1(x)(sa_1(x)a_1'(x) - a_1(x)a_1'(x) - sa_2'(x))$, which is zero when the denominator $D(x)$ is zero. Consequently the [expression \(6\)](#) reduces to

$$f(x) = \frac{a_1'(x)}{2}. \tag{7}$$

In this case $g(x)$ takes the form

$$g(x) = \frac{a_1(x)a_1'(x)((s-2)(s-1)a_1^2(x) + 2s^2a_2(x)) - 4s^2a_2(x)a_2'(x)}{2s^2((s-1)a_1^2(x) - 2sa_2(x))}. \tag{8}$$

In the above expression of $g(x)$ since the denominator divides the numerator, working in the same way as we did for the polynomial $f(x)$ we obtain that

$$g(x) = \frac{a_2'(x)}{s} - \frac{a_1(x)a_1'(x)}{2s},$$

and this completes the proof of the corollary. \square

We observe that the generalized Liénard polynomial differential system of [Example 1](#) does not satisfy the assumption that the common denominator of the expressions of f and g stated in [Theorem 1](#) vanishes. So we cannot apply [Corollary 2](#) to differential system (5).

Proof of Proposition 3. Under the assumptions of the proposition we arrive to an algebraic system between the parameters of $a_i(x)$ for

$i = 1, 2, 3$. The equation for the highest power in x which corresponds to x^{11} is $3a_{13}^2(4a_{13}a_{33} - a_{23}^2) = 0$.

First we consider $a_{33} = a_{23}^2/(4a_{13})$ with $a_{13} \neq 0$. Then vanishing the coefficients associated to the next powers of x we obtain $a_{32} = (2a_{13}a_{22}a_{23} - a_{12}a_{23}^2)/(4a_{13}^2)$, $a_{31} = (a_{13}^2a_{22}^2 + 2a_{13}^2a_{21}a_{23} - 2a_{12}a_{13}a_{22}a_{23} + a_{12}^2a_{23}^2 - a_{11}a_{13}a_{23}^2)/(4a_{13}^3)$, and

$$a_{30} = \frac{1}{8a_{13}^4}(4a_{13}^3a_{21}a_{22} - 2a_{12}a_{13}^2a_{22}^2 + 4a_{13}^3a_{20}a_{23} - 4a_{12}a_{13}^2a_{21}a_{23} + 4a_{12}^2a_{13}a_{22}a_{23} - 4a_{11}a_{13}^2a_{22}a_{23} - 2a_{12}^2a_{23}^2 + 4a_{11}a_{12}a_{13}a_{23}^2 - 2a_{10}a_{13}^2a_{23}^2 + a_{13}a_{33}^2).$$

But the solutions obtained vanishing the coefficients of next powers of x provides a reducible invariant algebraic curve $F = 0$.

Second we consider $a_{13} = 0$. The next coefficient is $2a_{12}^2a_{23}^2 = 0$. The case $a_{23} = 0$ with $a_{12} \neq 0$ implies $a_{23} = a_{33} = 0$ and $a_{32} = a_{22}^2/(4a_{12})$, and all the invariant curves $F = 0$ are reducible. The case $a_{12} = 0$ implies $a_{23} = 0$. Imposing that the rest of equations vanish we obtain the polynomials f, g and F of the statement of the proposition. \square

Proof of Proposition 4. This proof follows in a similar way to the proof of Proposition 3, but firstly imposing that f and g satisfy (7) and (8). We omit the details that the reader can compute easily. \square

CRedit authorship contribution statement

Jaume Giné: Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Writing – original draft, Writing – review & editing, Visualization. **Jaume Llibre:** Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Writing – original draft, Writing – review & editing, Visualization.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The author is partially supported by the Agencia Estatal de Investigación grant PID2020-113758GB-I00 and an AGAUR (Generalitat de Catalunya) grant number 2017SGR 1276. The second author is partially supported by the Agencia Estatal de Investigación grant PID2019-104658GB-I00, and the H2020 European Research Council grant MSCA-RISE-2017-777911.

References

- [1] Christopher CJ. Liouvillian first integrals of second order polynomial differential equations. *Electron J Differential Equations*. 1999;(49) 7 pp.
- [2] Dumortier F, Llibre J, Artés JC. *Qualitative theory of planar differential systems*. New York: Universitext, Springer-Verlag; 2006.
- [3] Singer MF. Liouvillian first integrals of differential equations. *Trans Am Math Soc*. 1992;333:673–88.
- [4] Zhang X. *Integrability of dynamical systems: algebra and analysis*. Developments in mathematics. Singapore: Springer; 2017.
- [5] Chavarriga J, García IA, Llibre J, Zoladek H. Invariant algebraic curves for the cubic Liénard system with linear damping. *Bull Sci Math*. 2006;130(5):428–41.
- [6] Cherkas LA. Liénard systems for quadratic systems with invariant algebraic curves. *Differ Uravn*. 2011;47(10):1421–7. translation in *Differ. Equ*. 47 (2011), no. 10, 1435–1441.
- [7] Giné J. A note on: the generalized Liénard polynomial differential systems $x = y, y = -g(x) - f(x)y$, with $\text{deg } f + 1, \text{ are not Liouvillian integrable}$. *Bull Sci Math*. 2015;139:214–27. *Bull. Sci. Math.* 161 (2020), 102857.
- [8] Giné J, Llibre J. Invariant algebraic curves of generalized Liénard polynomial differential systems. *Mathematics*. 2022;10(209) 5 pp.
- [9] Hayashi M. On polynomial Liénard systems which have invariant algebraic curves. *Funkc Ekvacioj*. 1996;39:403–8.
- [10] Hayashi M. A note on Liénard systems with invariant algebraic curves. *Adv Differ Equ Control Process*. 2010;6(1):15–24.

- [11] Llibre J, Valls C. The generalized Liénard polynomial differential systems $x' = y$, $y' = -g(x) - f(x)y$, with $\deg g = \deg f + 1$, are not liouvillian integrable. *Bull Sci Math.* 2015;139:214–27.
- [12] Zolaček H. Algebraic invariant curves for the Liénard equation. *Trans Am Math Soc.* 1998;350(4):1681–701.
- [13] Liu C, Chen G, Yang J. On the hyperelliptic limit cycles of Liénard systems. *Nonlinearity.* 2012;25:1601–11.
- [14] Qian X, Yang J. On the number of hyperelliptic limit cycles of Liénard systems. *Qual Theory Dyn Syst.* 2020.;19(43).
- [15] Qian X, Shen Y, Yang J. Invariant algebraic curves and hyperelliptic limit cycles of Liénard systems. *Qual Theory Dyn Syst.* 2021.;20(44).
- [16] Chèze G, Cluzeau T. On the nonexistence of liouvillian first integrals for generalized Liénard polynomial differential systems. *J Nonlinear Math Phys.* 2013;20(4):475–9.
- [17] Llibre J, Valls C. Liouvillian first integrals for Liénard polynomial differential systems. *Proc Am Math Soc.* 2010;138(9):3229–39.
- [18] Llibre J, Valls C. Liouvillian first integrals for generalized Liénard polynomial differential systems. *Adv Nonlinear Stud.* 2013;13(4):825–35.
- [19] Llibre J, Valls C. Liouvillian first integrals for a class of generalized Liénard polynomial differential systems. *Proc Roy Soc Edinburgh Sect A.* 2016;146(6):1195–210.
- [20] Valls C. Liouvillian integrability of some quadratic Liénard polynomial differential systems. *Rend Circ Mat Palermo.* 2019;68(3):499–519.
- [21] Demina MV. Invariant algebraic curves for Liénard dynamical systems revisited. *Appl Math Lett.* 2018;84:42–8.
- [22] Demina MV. Novel algebraic aspects of liouvillian integrability for two-dimensional polynomial dynamical systems. *Phys Lett A.* 2018;382(20):1353–60.
- [23] Demina MV, Giné J, Valls C. Puiseux integrability of differential equations. *Qual Theory Dyn Syst.* 2022;21(35):1–35.
- [24] Demina MV, Valls C. On the Poincaré problem and liouvillian integrability of quadratic Liénard differential equations. *Proc Roy Soc Edinburgh Sect A.* 2020;150(6):3231–51.
- [25] Llibre J, Valls C. On the local analytic integrability at the singular point of a class of Liénard analytic differential systems. *Proc Am Math Soc.* 2010;138(1):253–61.